



Biofabricated nanoparticles: Their delivery and utility in *Plutella xylostella* management

Kapinder^{1*}, Kriti Bhardwaj¹, Savita Singh¹ & Tarkeshwar²

¹Department of Zoology, Applied Entomology Lab, DDU Gorakhpur University, Gorakhpur-273 009, Uttar Pradesh, India

²Department of Zoology, Kalindi College, University of Delhi, Delhi-110 008, India

Received 22 December 2021; revised 13 February 2022

The diamondback moth (DBM), *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) is one of the most serious cosmopolitan pests of cruciferous plants. It causes severe damage to crop quality and production, with estimated global losses of 20-40% per year. Consumer preference regarding pest-free and uncontaminated crop, unavailability of highly competitive natural enemies as well as migratory nature of DBM has obligated for the mandatory use of insecticides. The over reliance and indiscriminate use of chemical insecticides to control DBM has lead to development of resistance against almost all the modern insecticides. In addition, it has also developed resistance against most of the *Bt* strain. To overcome the problem of resistance, there is an urgent need for new and highly efficient method to control DBM which must be eco-friendly, cost effective and has minimum impact on the environment. One of such alternatives is nanotechnology which fulfil all the criteria to become an ideal insecticide or delivery system for insect pest management especially DBM. Nanotechnology offers great improvement in the field of pesticides as it is less toxic, shelf-life enhancement and higher water solubility. Loading of Indoxacarb on nanoparticles are reported to be highly effective they suppressed the activity of detoxification enzymes such as GST, CarE, and P450. Green synthesis of AgNPs, Single walled carbon nano tube (SWCNT) also exhibits larvicidal and pupicidal effect. These are also reported to adversely affect food consumption, growth, pupation and fecundity of *P. xylostella*. These nanoparticles have been found to be more stable, also the controlled release of active ingredient for a long time and provides target specific control of *P. xylostella* for solving adverse situations of the crop fields like less food security, lesser food productivity and environmental imbalance.

Keywords: Biofabrication, Cruciferous plant, Nanoparticles, Nanopesticides, *Plutella xylostella*

Worldwide *Plutella xylostella* caused an enormous crop loss of 90% yields and annual damage upto \$4-5 billion per year. Management of the pest involves the cost of US \$1.0 billion per year¹. The conventional procedure like IPM used in crop fields are inadequate and utilization of huge amounts of chemical pesticides have negative effects on human beings and animals apart from the reduce in fertility of soil². Therefore, without harming the nature, nanotechnology has provided green and effectual liberty for crop protection in agriculture. Nanobiotechnology has been provided options to overcome pesticides, like decreasing toxicity, shelf-life enhancement and by making less water-soluble pesticides to water-soluble pesticides, even more of the positive environmental impacts. Nanotechnology has great involvement in the development of opportunities to deal with complex technical and economic issues of insect pest

management in agriculture field. Nanotechnology has developed in the last decade and has been able to create multiple creative materials with a huge range of forthcoming applications³. Nanoscale structures like nanomaterials and agrochemical entrapment of pesticides, fertilizers, herbicides, regulatory components of plant growth via involvement of surfactants, surface ionic attachments, dendrimers, polymers and other related components can be utilized in plant protection inputs using rapid delivery; it will ensure the rate of uptake which would be lower for active ingredients and directly reduces the quantity of agrochemical application by reduction of the input⁴ (Fig. 1). In terms of environment, the degradation prominence of nanotechnology delivered systems in crop fields is due to solubility and stability improvements⁵⁻⁸. Certain corporate companies have already been established in the marketing field to produce nano-scale emulsions in the form of microencapsulated pesticides because of the environmental trigger, loading of nanoparticles

*Correspondence:
E-mail: kapsind2007@gmail.com




Fig. 1 — Use of nanoparticles for the management of *P. xylostella*

entrapment with pesticides (with slower release in the field), low dose of nanocides products found to be effective in agricultural field protection and their lofty reactivity at nanoscale compared to their mass counterparts⁹.

A lepidoptera insect pest of cruciferous plants, *P. xylostella* (Lepidoptera: Plutellidae) is a cosmopolitan and notorious pest worldwide. In the fields, this pest causes 90% of the total yield loss. DBM larvae feed voraciously on leaves of cruciferous plants and causes major damage upto the level of foliar tissue of the leaf veins. The damage of larval stages of diamond back moth (DBM) could be seen in the plants without seedlings and head disruption in broccoli, cabbage, and cauliflower. The florets infested with larvae leads to complete deterioration of product upto the level of plant tissue removal. Diamondback moth was long considered a relatively insignificant minor pest. Its impact was overshadowed by such serious defoliators as imported cabbageworm. Many control measures are accepted for the management of this pest; whereas the most powerful technique to control the damage has involved the heavy dose of chemical insecticides¹. Overuse of synthetic chemicals proved to cause adverse effects such as insecticidal resistance, environmental imbalance due to their long persistence in the surroundings. According to the present scenario, the urgency evolved to develop an alternative method for control of this insect pest simultaneously this method

should be environmental friendly, more effective and provide safer prospective for the economy. The feasible alternative and evolving methods included formation of biofabricated nanoparticles with the help of plant or the plant parts either in the form of extracts or any active compounds for the protection of the field crops¹⁰. Extra cts of several plants such as bitter gourd, clove, neem, bakain, eucalyptus, datura, ginger and garlic were used for biofabrication of silver nanoparticles and found very effective against *P. xylostella* larvae. The present review has focused on two main aspects of nanoparticles where NPs were synthesized and utilized for management of insect pest *P. xylostella*, either in the form of nanoparticles itself or as carriers for the targeted delivery system and protectants entrapping fungicides, insecticides and herbicides.

Insecticidal activity of nanoparticles

Silicon dioxide nanoparticles (SiO₂NPs)

The nanosilica exhibited enormous entomotoxic effects on the *P. Xylostella* larvae. Biologically derived nano-silica from wild sugarcane, *Erianthus arundinaceus* showed insecticidal activity against larvae of diamond back moth, *P. xylostella* with LC₅₀-30.03 µg/mL¹¹. The nanosilica prepared through sol-gel technique and mortality of insect larvae was demonstrated by spray methods, leaf dipping, larva dipping and dust spray methods. The highest mortality of larvae was reported in dust spray method

where mortality was increased from 58% to 85% at 24 h and 72 h of treatment using 1 mg/cm² concentration. This mortality rate could be further increased with increasing concentrations and duration of exposure. The death of feeding larvae occurred due to desiccation, abrasion of body walls and spiracle blockage¹².

Single walled carbonnanotube (SWCNT)

It was produced by catalytic vapour deposition (CVD) and reported to exhibit mortality in *P. xylostella* larvae only at higher concentrations such as 138.24 µg/mL concentration, larval feeding reduced upto 10% and the morphology of the larvae also demonstrated darker colour which implies that SWCNT has been deposited in the larval gut. As concentration of carbon nanotubes increased there was significant reduction in pupation rate, as well as fecundity (egg/adult) rate also declined upto 45% with increasing SWCNT concentrations. The SWCNT feeding larvae showed no or minor increase in larval size with increasing food consumption¹³.

Titanium dioxide nanoparticles (TiO₂NPs)

Titanium dioxide nanoparticles (TiO₂) in its ionic form has been considered as active compound and proved as available alternative to the synthetic pesticidal control in the management of pest in the crop fields. Titanium dioxide TiO₂-NPs synthesised by sol-gel method¹⁴ and dose dependant response was obtained through regression analysis for the bioassay of titanium dioxide (TiO₂) while tested on diamondback moth larvae with different concentration and different time period interval, respectively. The cumulative mortality was nearly 50%. The above result was supported by Kookana *et al.* (2014)¹⁵ who suggested that nanomaterials could be an alternative approach against synthetic insecticides for insect pest management.

Emamectin benzoate nanoformulations

This approach utilizes colloidal form of delivery system which effectively controls pesticides delivery to increase photostability and efficacy of semi-synthetic and natural pesticides. The prepared emamectin benzoate nanoformulations involved polymeric nanocapsules and nanosilica of two types, mesoporous and silicon dioxide which worked as carriers for emamectin benzoate. This type of combination displayed high encapsulation efficacy of upto 70-90% in different nanocapsules. The efficacy

of nanocapsules against *P. xylostella* was found to be EB+SNPs had more effective in comparison to other EB+NFs and EB placebo. The LC₅₀ values for EB+SNPs, EB+MCM-48, EB+PNC and EB placebo would be 0.18, 4.03, 8.49 and 11.06 mg, respectively¹². The enhanced efficacy of nanocapsulations was due to smaller particle size, large surface area and their mobility increment ratios which lead to the high penetrance ratio in the larval body causing higher mortality at smaller dosage.

Silica nanoparticles (SiNPs)

Because of the known properties of the controlled release, the major attention had been given to the silica nanoparticles for smart delivery system potentials. For the preparation of florescent mesoporous silica NPs with embedding of Carbon dots (FL-SiO₂ NPs), silica precursors had been used *via* tetraethoxysilane (TEOS) involvement and structure-directing agents as Cetyltrimethylammonium bromide (CTAB). The leaf-dipping method was used for the detection of insecticidal activity against insect larvae. In this experimental setup the control has been Indoxacarb technical concentration (TC) and various gradients of descending order 25, 12.5, 6.25, 3.12, and 1.56 mg/L had been tested. The mortality of larvae after 48 h of treatment was 90% for IN@FL-SiO₂ NPs and for control data 76% at 25 mg/mL of higher concentration. The FL-SiO₂ NPs presence had been observed at the midgut of *P. xylostella* using confocal laser scanning microscopy (CLSM)¹⁶. The higher mortality of *P. xylostella* larvae in the treatment was observed due to decreased enzymatic activities which could imply the nano-based delivery system of insecticides for the decreasing or delaying the effect of insecticide resistance. This method could be included for resistance selection of pest to IN@FL-SiO₂ NPs and for identification of resistant strains of diamondback moth using bioactive and biochemical assay¹⁷.

The effect of microsizechlorfenapyr (MCSC) and nanosized chlorfenapyr (NCSC) against *P. xylostella* was analysed¹⁸ and found that NCSC caused higher mortality in the larvae and also required half dose of treatment as compared to MCSE. The higher larval mortality was observed from first day and subsequent days that confirmed the reduced amount of pesticides as well as less time interval through applying nano silicons. The enhanced activity had been reported due to improvement in biological efficacy of chlorfenapyr-loaded silica as well as silica itself provided insecticidal activity¹⁸. Kaziem *et al.* (2018)¹⁹

reported that avermectin (AVM) controlled released using α -Amylase carriers had been prepared using lair mesoporous silica with α -CD as capping molecule. The AVM-CRF biological tests were conducted on *P. xylostella* larvae through uncapping of α -CD enzyme which later on release AVM causing larval deaths. The combination had been able to keep 0.6 mg/L AVM to be in bull form for 14th of the subsequent days showing 83.33% mortality of the insect larvae.

Biofabrication of nanopesticides

A lot of techniques have been designed using variety of physical, chemical methods for synthesis of nanomaterials. This field had shown different properties of the same chemical in the bulk scale and in the nano scale. In the nanoscale, because of larger surface area the atoms provided more chemical reactions and the time of exposure also increases to initiate any reaction. As we have seen in the above context, with the help of nanotechnology variety of new pesticides, insecticides and insect repellents had been prepared. Some chemicals used during the synthesis of these nanoparticles in polyol method, microemulsion, electrochemical decomposition, and thermal decomposition are found to be very toxic for the environment as well as for human beings. To overcome this negative effect of the NPs we are on the brink to developed methods of preparation involving green synthesis approach. The new emerging arena of green synthesis involves plants

metabolites for the reducing agents, oxidising agents, and so on. These plant products used for biofabrication are cost-effective, environment friendly and with minimum adverse effects of the human beings⁴. The entrapment of pesticides inside polymers increases the stability of the compound and causes slow release which finally decreases the dose of the same compound and also decreased side effects. There are still a lot of unexplored fields available to be worked on. The list given below provides the successful biosynthesis of NPs used in the agricultural field to control the insect pest *P. xylostella* (Table 1).

Biopolymer based silver nanoparticles (AgNPs)

Biofabrication of silver nanoparticles (AgNPs) have been considered as highly reliable method²¹ to use it as an alternative way to manage the population of *P. xylostella*. Biosynthesized silver nanoparticles (AgNPs) using neem plant demonstrated highest mortality of 76% at a concentration of 2.4 mg/mL and minimum mortality of 28% at the concentration of 0.15 mg/mL against the 3rd instar larva of diamondback moth for the 24 h of time exposure. Proceeding of the same experiment for 48 h showed that 80% and 32% of mortality for the maximum and minimum concentrations, respectively. The result for 72 h found to be 92% and 36% of the mortality for the higher and lower concentrations, respectively¹.

When clove (*Syzygium aromaticum*) was used for biofabrication of AgNPs and its effect was evaluated on larvae of diamond back moth for 24 h, larval

Table 1- Application of various nanoparticles for the management of *Plutella xylostella*

Nanoparticles	Synthesis	LC ₅₀	Mode of action	References
Nano-silica	sugarcane	30.03 μ g	-	Kannan <i>et al.</i> 2017
Silicon dioxide	Sol-gel technique	0.99 mg/L	Dessication, abrasion of body walls	Shoaib <i>et al.</i> 2018
SWCNT	Catalytic vapour deposition	138.28 μ g/ mL	Stored in the gut with zero metabolism and bind with actin filament centromere	Taiaba Afrin 2017
Titanium dioxide	Sol-gel method	Cumulative mortality	-	Kokkan <i>et al.</i> 2014
Emamectin benzoate nanoformulation	Colloidal form for delivery	0.18 mg	Penetration into the larval body	Shoaib <i>et al.</i> 2018
Silica nanoparticles AVM-CRF	Chemical method	0.6 mg/L	Slower release of pesticide AVM	Kaziem <i>et al.</i> 2018
Silver nanoparticles	<i>H. musciformis</i>	24.5 ppm	Decreases fecundity and longevity	Roni <i>et al.</i> 2014
Neem oil	Neem extract biosynthesis	0.94 mg/ mL	Slower release of neem oil	Forim <i>et al.</i> 2013
Bt-based nanopesticides	Top-down approach homogenization using water	-	Crystal toxin in the midgut alkaline medium	Kim and Joe, 2012

mortality of 76% at the dose of 2.8 mg/mL and 24% for the dose of 0.175 mg/mL was observed. After 48 h, the mortality was further increased to 84% and 32%, respectively. After the treatment for 72 h the mortality was 88% and 36% for the higher dose and lower dose where the highest dose mortality had been 80% and 88% whereas the lowest dose of mortality of 32% and 36%¹. Synthesis of AgNPs using bitter gourd (*Momordica charantia*) added further toxicity when tested against *P. xylostella* larva for the maximum concentration of 3 mg/mL and minimum concentration of 0.1875 mg/mL showed 72% and 24% of mortality, respectively. After 72 h of treatment, the reported mortality for the same concentration found to be 84% and 36%, respectively¹.

Biofabrication of AgNPs using *Datura* at a concentration of 2 mg/mL exhibited highest mortality of 72% and for the minimum dose of 0.125 mg/mL reported 28% of effects for only 24 h of time duration. After treatment of 48 h, larval mortality increased upto 80% and 32% for higher and lower concentration, respectively, and the report of 72 h recorded values of 84% and 40% for the former and later, respectively¹. AgNPs biosynthesized using *H. musciformis* sea weed leaf extract exhibited larvicidal and pupicidal toxicity at low concentration against *P. xylostella* and also decreased fecundity and longevity of adults of diamondback moth. The LC₅₀ value of AgNPs for L₁ and pupae had been reported as 24.5 ppm and 38.23 ppm for *P. xylostella*²⁰.

Silver nanoparticles biosynthesis using bakain (*Melia azedarach*) was found to exhibit toxicity against insect larvae because of the toxic effect of the plant itself¹. When these Ag-NPs were tested against *P. xylostella* for 24 h showed the maximum mortality of around 72% for the dose of 4 mg/mL and minimum mortality rate of around 28% for the dose of 0.25 mg/mL when the experiment proceeded for 72 h at the dose of 0.1875 mg/mL showed 44% of larval mortality and the maximum dose of the same experiment would be recorded at 3 mg/mL of 80% mortality implies killing of 20 out of 25 larvae in the experimental apparatus¹. AgNPs biofabrication using garlic demonstrated highest mortality of *P. xylostella* larvae of 72% after 24 h of treatment²¹. When this preceded further for 24 h on the subsequent day exhibited 80% of the total mortality. When further proceeding upto 72 h, the larval mortality was recorded to be 84%²¹. It had also been reported that garlic also exhibited insecticidal activity against *P. xylostella* larvae²².

Nanocapsules

Biofabrication of nanocapsules were performed using *Azadiracta indica* oil and neem extracts involving polymer and emulsion had been tested against *P. xylostella* larvae. Kale leaves dipped in different concentrations of treatment were provided as food for the larvae. The result exhibited 100% mortality in first instar larvae on the 9th day of treatment as well as powdered NPs caused only 91.7% mortality²³. Same results were also demonstrated by Torres *et al.* (2006)²⁴ where neem extract caused 100% larval mortality. The different mechanisms of action involved in the solution phase dipped with kale leaves included solvent phases causes active compound release earlier whereas the powdered form caused later release of active components, greatly dependent on the colloidal silica hydration. This provided a bio-sustainable process involved no harmful chemicals.

Bt-based nanopesticides

This category includes Bt size decreased via Bt technical powders using top-down method as well as Bt-homogenization through water as an emerging method. Smaller sized Bt-homogenized particles caused increased DBM larval mortality. The unhomogenized powder caused larval mortality of 27.5% within 2 days of treatment which increased upto 72.6% in day three whereas homogenized Bt-nanopesticides exhibited 78.3% even on day two and 100% mortality on day three²⁵. The mechanism of action involved in nanoparticles of Bt enhanced the solubility of the crystal toxin in the midgut alkaline media which caused more production of small size toxins that actively binds with the receptors and ultimately caused rapid gut paralysis followed by larval death.

Conclusion

The nanotechnology has provided promising methods for the successful implementation in the field of agriculture for the development of nanopesticides. The green synthesis approach uses plants or plant products which significantly decreased our reliance over the use of toxic chemicals. The biofabrication of silver using various plant materials such as garlic, cloves, bitter gourd, bakain, which is more effective than its simple NPs synthesized using chemicals. The plants products itself also have pesticidal activity which further improves the efficacy of nanopesticides and subsequently decreases the dosage of that pesticides. So, it has been concluded that the

biosynthesis using green approach would be a significant step towards the development of nanotechnology with minimum or no less at all.

Acknowledgement

We are thankful to Deen Dayal Upadhyaya Gorakhpur University for providing the facility to complete the present task.

Conflict of interest

All authors declare no conflict of interest.

References

- 1 Ali F, Tariq M & Shaheen FA, Toxicity of different plant extracts and green silver nanoparticles against *Plutella xylostella* (Lepidoptera Plutellidae). *Plant Protection*, 3 (2019) 251.
- 2 Goswami A, Roy I, Sengupta S & Debnath N, Novel applications of solid and liquid formulations of nanoparticles against insect pests and pathogens. *Thin Solid Films*, 519 (2010) 1252.
- 3 Bhattacharyya A, Bhaumik A, Usha Rani P, Mandal S, & Eparti TE, Nano-particles - A recent approach to insect pest control. *Afr J Biotech*, 9 (2010) 3489.
- 4 Kapinder, Dangi K & Verma AK, Efficient & eco-friendly smart nano-pesticides: Emerging prospects for Agriculture. *Mater Today: Proceed*, 45 (2021) 3819.
- 5 Chen H & Yada R, Nanotechnologies in agriculture: newtools for sustainable development. *Trend Food Sci Technol*, 22 (2011) 585.
- 6 Johnston CT, Probing the nanoscale architecture of clay minerals. *Clay Minerals*, 45 (2010) 245.
- 7 Jha Z, Behar N, Sharma SN, Chandel G, Sharma D & Pandey M, Nanotechnology: prospects of agricultural advancement. *Nano Vision*, 1 (2011) 88.
- 8 Sastry RK, Rashmi H, Rao N & Ilyas S, Integrating nanotechnology into agri-food systems research in India: a conceptual framework. *Technol Forecast Soc Change*, 77 (2010) 639.
- 9 Debnath N, Das S, Seth D, Bhattacharya RC & Goswami A, Entomotoxic effect of silica nanoparticles against *Sitophilus oryzae* (L.). *J Pest Sci*, 84 (2011) 99.
- 10 Mari JM, Efficacy of different plant extracts against diamondback moth, *Plutella xylostella* (L.) on cauliflower. In *Int Conf Appl Life Sci*, (IntechOpen) 2012.
- 11 Kannan M, Bhaskaran A, Prithiva JN, Preetha S & Lokesh S, Biogenic silica nanoparticles from *Erianthus arundinaceus* as novel insecticide against diamond back moth, *Plutella xylostella* L. Proceedings of the International Symposium on Sugarcane. Hotel Le Meridian, Coimbatore: Sugarcane Breeding Institute and Tamil Nadu Agricultural University, Coimbatore (2017) 224.
- 12 Shoaib A, Elabasy A, Waqas M, Lin L, Cheng X, Zhang Q & Shi ZH, Entomotoxic effect of silicon dioxide nanoparticles on *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) under laboratory conditions. *Toxicol Environ Chem*, 100 (2018) 80.
- 13 Afrin T, Effects of engineered carbon and silver nanoparticles on *Plutella xylostella* consumption, growth, population, survival and fecundity. MSU Graduate Thesis (2017).
- 14 Preetha S, Kannan M, Lokesh S, Viji N, Prithiva JN & Gowtham V, Titanium dioxide (TiO₂) nanoparticles as a novel insecticide against diamondback moth, *Plutella xylostella* L. in cauliflower. *Trends in Biosciences*, 11 (2018) 2999.
- 15 Kookana RS, Boxall ABA & Reeves PT. Nanopesticides: Guiding Principles for Regulatory Evaluation of Environmental Risks. *J Agric Food Chem*, 62 (2014) 739.
- 16 Bilal M, Xu C, Cao L, Zhao P, Cao C, Li F & Huang Q, Indoxacarb-loaded fluorescent mesoporous silica nanoparticles for effective control of *Plutella xylostella* L. with decreased detoxification enzymes activities, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, No. 2 Yuanmingyuan West Road, Haidian District, Beijing 100193, P. R. China (2020).
- 17 Benelli G, Mode of action of nanoparticles against insects. *Environ Sci Poll R*, 25 (2018) 12329.
- 18 Song MR, Cui SM, Gao F, Liu YR, Fan CL, Lei TQ & Liu DC, Dispersible silica nanoparticles as carrier for enhanced bioactivity of chlorfenapyr. *J Pest Sci*, 37 (2012) 258.
- 19 Kaziem AE, Gao Y, Zhang Y, Qin X, Xiao Y, Zhang Y, You H, Li J & He S, α -Amylase triggered carriers based on cyclodextrin anchored hollow mesoporous silica for enhancing insecticidal activity of avermectin against *Plutella xylostella*. *J Hazard Mater*, 359 (2018) 213.
- 20 Roni M, Murugan K, Panneerselvam C, Subramaniam J, Nicoletti M, Madhiyazhagan P, Dinesh D, Suresh U, Khater HF, Wei H, Canale A, Alarfaj AA, Munusamy, MA, Higuchi A & Benelli G, Characterization and biotoxicity of *Hypneamusiformis*-synthesized silver nanoparticles as potential eco-friendly control tool against *Aedes aegypti* and *Plutella xylostella*. *Ecotoxicol Environ Safety*, 121 (2015) 31.
- 21 Sarker P, Rahman M & Das B, Effect of intercropping with mustard with onion and garlic on aphid population and yield. *J Bio-Sci*, 15 (2007) 35.
- 22 Reuben SO, Yahya SN, Misangu RN & Mulungu LS, Field evaluation on effects of common spices in the control of diamondback moth (*Plutella xylostella* L.) pest of Chinese cabbage (*Brassica campestris* L.) commercial cultivar. *Asian J Plant Sci*, 5 (2006) 85.
- 23 Forim MR, Costa ES, Da Silva MFDGF, Fernandes JB, Mondego JM & Boica Junior AL, Development of a new method to prepare nano-microparticles loaded with extracts of *Azadirachta indica*, their characterization and use in controlling *Plutella xylostella*. *J Agri Food Chem*, 61 (2013) 9131.
- 24 Torres AL, Júnior BAL, Medeiros CAM & Barros R, Effect of aqueous extracts of *Azadirachta indica*, *Melia azedarach* and *Aspidosperma pyrifolium* on the development and oviposition of *Plutella xylostella*. *Bragantia*, 65 (2006) 447.
- 25 Kim JS & Je YH, Milling effect on the control efficacy of spray dried *Bacillus thuringiensis* technical powder against diamond back moths. *Pest Manag Sci*, 68 (2012) 321.