



Utilization of fungal biocontrol agents against rice sheath blight disease provides insight into their role in plant defense responses

Bishnu Maya Bashyal^{1*}, Shashi Pandey^{1,2}, Ayush Raj Singh^{1,2}, Prashantha ST¹, Gopalakrishnan S², Dinesh Singh¹,
Deeba Kamil¹ & Rashmi Aggarwal¹

¹Division of Plant Pathology; &²Division of Genetics, ICAR-Indian Agricultural Research Institute,
New Delhi-110 012, Delhi, India

Received 26 September 2022; revised 19 October 2022

Biotic and abiotic factors have an effect on rice production all around the world. Diseases are regarded as major restrictions among the biotic stressors, and rice sheath blight (*Rhizoctonia solani* Kühn) is one of the most calamitous diseases that significantly damage the crop. Lately, biocontrol of fungal plant pathogens has appeared as an appealing approach. The present investigation was undertaken to evaluate different biocontrol agents like *Talaromyces flavus*, *Chaetomium globosum*, *Pseudomonas fluorescens* and *Aspergillus niger* against sheath blight disease. Prior to sowing, seeds were bioprimered with each isolate and sown in the nursery. After 21 days, seedlings were transplanted *in-vivo* and were inoculated with a virulent isolate of *Rhizoctonia solani* at maximum tillering stage. Observations on biochemical parameters and gene expression studies were carried out at 24, 48, 72, and 96 hpi. Enzymatic activity *viz.*, chitinase, β -1,3-glucanase, catalase, and PAL was observed maximum in *Chaetomium globosum*. PR-genes *viz.*, IPT, BrD, HmPr, AMP, AldD, NIC and LisH showed up-regulation at 96 hpi. *Chaetomium globosum* had the highest yield, maximum number of tillers with least relative lesion height (RLH%) as compared to other treatments. However, results indicated biocontrol agents are helpful and they induce multitude of defence responses against *R. solani* in rice.

Keywords: Biocontrol agents, PAL, PB-1121, *Rhizoctonia solani*, Sheath blight

Rice, the world's most significant food crop, has been produced throughout South Asia for over 6000 years. Rice is currently the staple diet of about four billion people, or half of the world's population¹. Various biotic and abiotic factors have an impact on rice production around the world. Diseases are one of the most important biotic pressures on rice production, with rice sheath blight, caused by *Rhizoctonia solani* Kühn, being one of the most serious diseases capable of causing major crop damage in areas where intensive agriculture is practised². This disease is also called as "snake skin disease," "mosaic foot stalk," and "rotten foot stalk" because to its distinctive symptoms³⁻⁴. Rice sheath blight is a serious damaging disease that causes quality degradation and up to a 50% of yield loss each year around the world⁵⁻⁷. In India, annual losses due to sheath blight (ShB) are estimated to be 10%⁸. For the management of this disease, chemical control strategies are already in use, but a strategy for eco-friendly compounds with enhanced disease control potential is critical⁹.

Rhizoctonia solani is a phytopathogenic fungus that infects a wide variety of plants¹⁰. This disease affects a lot of plant groups, including the Asteraceae, Brassicaceae, Fabaceae, Poaceae, and Solanaceae, as well as several woody ornamental plants and forest trees¹¹⁻¹⁴. Biological management of fungal plant diseases has recently emerged as an appealing and feasible option which helps disease management as well as improves the plant growth characters also. The principal antagonistic microorganisms against numerous plant pathogens are *Chaetomium* spp., *Aspergillus* spp., *Pseudomonas* spp., and *Talaromyces* spp., which were used as biological control for the pathogens such as *Gaeumannomyces graminis* var. *tritici*¹⁵, *Fusarium oxysporum*¹⁶, *Sclerotium rolfsii*¹⁷, *Verticillium dahliae*¹⁷, *Verticillium albo-atrum*¹⁸, *Rhizoctonia solani*¹⁹ and *Sclerotinia sclerotiorum*²⁰⁻²¹. *Chaetomium globosum* has the potential to be a biocontrol agent for a variety of seed and soil-borne diseases²². In general, bio-control agents work through competition, mycoparasitism, antibiosis, or a combination of these mechanisms to stop the growth of bacteria and fungus²³. *Pseudomonas fluorescens* has emerged as a promising biocontrol agent for plant

*Correspondence:
E-mail: bishnumayabashyal@gmail.com

disease suppression by protecting seeds and roots against fungal infection. They have been shown to promote plant development and lower the severity of numerous fungal infections²⁴⁻²⁵. *Aspergillus niger* has an antagonistic relationship with *R. solani*. It lowers the occurrence of disease²⁶.

Plant defence induction is a novel biological strategy for controlling plant disease. Many biocontrol agents are being studied for their ability to induce plant systemic resistance to pathogenic fungus. There are definite signs of systemic defense-related enzyme activity in the soybean, tomato, and *Arabidopsis thaliana* plants. These enzymes include phenylalanine ammonia lyase (PAL)²⁷⁻³⁰.

Most hyperparasitic biocontrol agents produce enzymes like chitinases and glucanases. Plant pathogens can use specific methods to defend themselves against cell wall destroying enzymes. In fact, *R. solani* secretes a wide variety of effectors, enzymes that are active on carbohydrates, and various kinds of secondary metabolites that have an impact on host immunity³¹⁻³³. Plants have numerous layers of defences against pathogen attacks to counterbalance the effects of pathogenicity factors in plant diseases. Pathogenicity-related proteins such as β -1,3-glucanase, and chitinase plays a critical role in fungal-pathogen infection defence³⁴. β -1,3-glucanase hydrolyzes β -1,3-glucans, which are found in most fungal cell walls, whereas chitin is a significant component of fungal cell walls that is hydrolyzed by the chitinase enzyme³⁵.

Here, we tested four different biocontrol agents: *Chaetomium globosum* (Cg 2), *Pseudomonas fluorescens* (DTPF-3), *Aspergillus niger* (ITCC no. 7790), and *Talaromyces flavus* (Tf2) against *Rhizoctonia solani* inoculated plants to see how they affected overall plant growth and development, as well as defence response to the phytopathogenic fungus *R. solani*. As a result, numerous plant growth parameters were examined in the presence of pathogen and biocontrol agents. In addition, the activities of defense-related enzymes were characterised, and the expression of distinct pathogenicity-related genes in plants was studied.

Materials and Methods

Location of experiment

All the experimental studies were carried out at ICAR-Indian Agricultural Research Institute (IARI), Pusa campus, New Delhi at latitude of 28.08°N and longitude of 77.12°E, the height above mean sea level being 228.61 metres (750 feet).

Fungal isolates, rice genotypes taken for the study

Highly virulent isolate of *R. solani* collected from various rice growing areas in Uttar Pradesh, India. R-359 (ON383512) was selected for inoculation. Slants of Potato Dextrose Agar (PDA) were used to culture and preserve isolates, and they were kept at 4°C. For present study, the susceptible rice cultivar Pusa Basmati-1121 was obtained from Division of Genetics, IARI, New Delhi.

Biocontrol agents and seed treatment

Biocontrol agents *Talaromyces flavus* Tf2³⁶, *Chaetomium globosum* Cg2³⁷, *Aspergillus niger*³⁸, and *Pseudomonas fluorescens* DTPF-3³⁹ were taken from Division of Plant Pathology, ICAR-Indian Agricultural Research Institute, New Delhi. Spore suspensions of biocontrol agents prepared separately and passed through double-layered muslin fabric and concentration of spores were adjusted to 1×10^7 spores mL⁻¹ (*Talaromyces flavus*, *Chaetomium globosum* and *Aspergillus niger*) and 1×10^8 spores mL⁻¹ (*Pseudomonas fluorescens*). Seeds were dipped in biocontrol suspension for 24 h. Treated seeds were further sown in nursery bed prepared in glasshouse. Twenty-one days old nursery of the "Pusa Basmati 1121" treated with different biocontrol agents were uprooted carefully and transplanted at experimental field of ICAR-IARI. The normal agronomical practices with recommended doses of fertilizers (N: P₂O₅: K₂O: 120: 60: 60 kg ha⁻¹) were followed. Plots were kept weeds free through manual weeding. Seeds soaked only in water were treated as control.

Mass multiplication and inoculation of *Rhizoctonia solani*

Forty five days old transplanted plants (Maximum tillering stage) were inoculated with virulent isolate of *Rhizoctonia solani*. For which Typha (*Typha angustifolia*) stems were cut into 4-5 cm lengths, lightly filled 3/4th of a conical flask (500 mL), after that it was autoclaved for 15 min at temperature of 121°C, 15psi pressure. The Typha pieces were inoculated with fresh R-359 (ON383512) culture in a laminar air flow hood and incubated for one week at 26±1°C in BOD. As an inoculum, Typha pieces fully colonized with *Rhizoctonia solani* were utilised⁴⁰ in the rice genotype PB 1121. Inoculum (3 to 4 Typha bits) was inserted at the base of the tillers, and thread was used to bind the typha bits to the tillers, making contact between both the mycelium and rice plants. Three duplicates were kept for each isolates, as well as a control (three inoculated hills per pot were

considered as one replication and 3 replicates per isolates were evaluated).

Phenotypic evaluation of rice genotypes in field

Different parameters such as plant height, lesion height, numbers of tillers and grain yield were recorded for sheath blight disease severity. The relative lesion height (RLH) was calculated by using the following formula:

$$\text{RLH} = \frac{\text{Lesion height}}{\text{Plant height}} \times 100$$

Sample collection

For biochemical and real time PCR based analysis infected rice plants were collected at 24, 48, 72, and 96 h post inoculation. These samples were taken in triplicate and transferred to liquid N₂ after being covered in aluminium foil. After that, the samples were kept at -80°C for future examination.

Biochemical analysis

Enzyme extract preparation

In liquid N₂, one gram of each sample was crushed to a fine powder. At 4°C, 3 mL of sodium acetate buffer (50 mM, pH 5.2) was mixed with this tissue powder. At 4°C, the enzyme-buffer combination was centrifuged for 10 min at 10000 rpm. The supernatant was placed into a sterile tube and utilised in enzyme detection experiments.

Chitinase activity assay

The reduction of 3, 5-dinitrosalicylic acid in the presence of the amino-sugar N-acetyl-D-glucosamine (NAG) was used to determine chitinase enzyme activity. Chitin enzymatic hydrolysis releases NAG. The protocol for assaying chitinase was modified somewhat from that published by⁴¹.

In 1 mL of enzyme extract, we added 1 mL of 10% wet colloidal chitin (in 0.2 M sodium phosphate buffer, pH 6.5). The reaction mixture was then incubated at 50°C for 30 min. After that, we added 1 mL of 1% NaOH to the reaction mixture. To stop the enzymatic action, NaOH is utilized. The mixture was then centrifuged at 4°C for 10 min at 7500 rpm. In a sterile tube, the supernatant was collected. One mL of 1 percent 3, 5-dinitrosalicylic acid was combined with 1 mL of supernatant. 3, 5-dinitrosalicylic acid was dissolved in 30 percent sodium potassium tartrate and 2 M NaOH to make a solution. For 5 min, these tubes were immersed in boiling water. The reaction mixture was then diluted

1:10 with water after the tubes were allowed to cool to room temperature. At 535 nm, the absorbance of chitinase activity was measured. The amount of enzyme necessary to create 1 μmol of NAG in 1h was defined as a chitinase activity unit.

Four grams of chitin (Himedia) was added to 60 mL concentrated hydrochloric acid and stirred continuously at 4°C to make a colloidal chitin suspension. It was left at 4°C overnight after mixing. After 24 h, 500 mL of ice cold 95 percent ethanol was added without shaking and incubated at 4°C overnight. Chitin is precipitated by ice cold 95 percent ethanol. The ethanol was then removed by centrifuging the mixture at 10000 g for 10 min at 4°C. To remove the remaining ethanol, 10 mL ice cold sterile water was added to wash and centrifuged at 10000 g for 10 min at 4°C temperature. The washing process was repeated twice more before being autoclaved.

β-1,3-Glucanase activity assay

Abeles and Forrence's approach was used to detect β-1,3-glucanase activity. Laminarin was utilised as the substrate, and dinitrosalicylic was employed as the reagent, to measure the synthesis of reduced sugar⁴². In a sterile tube, 1 mL crude enzyme was added to 1 mL 2 percent laminarin solution. Tubes were incubated at 50°C for 60 min. Heating was used to incorporate laminarin into the reaction mixture. 3 mL dinitrosalicylic reagent was added to the reaction mixture after it had been heated. The tubes were heated at 100°C for another 5 min. The dinitrosalicylic reagent is used to halt the process. One gram NaOH, 0.2 g phenol, 0.05 g sodium sulphite, 18.2 g sodium potassium tartarate (Rochelle salt), and 1 g 3,5-dinitrosalicylic acid were combined in a 100 mL distilled water solution to make the dinitrosalicylic reagent. The reaction mixture was then diluted 1:10 with water after the tubes were allowed to cool to room temperature. At 500 nm, the absorbance was measured.

Phenylalanine Ammonia Lyase (PAL) activity assay

The enzyme activity of phenylalanine ammonia lyase (PAL) was measured using Mori *et al.*⁴³ technique. At 37°C, L-phenylalanine transforms to cinnamic acid. 0.4 mL Tris-HCl buffer (100 mM, pH 8.8) and 0.2 mL phenylalanine (40 mM) were added to 0.2 mL enzyme extract to make the reaction mixture. In place of enzyme extract, 30 μL extraction buffer (50 mM sodium acetate buffer, pH5.2) was employed as a control. The absorbance of the samples

was measured at 290 nm, with readings obtained every 30 sec for 5 min after the reaction began. PAL activity was measured in millimoles of cinnamic acid per milligrams of protein.

Catalase activity assay

The activity of the catalase enzyme was measured using the Bailly *et al.* technique⁴⁴. In the tube, we added 40 μ L of crude enzyme and 3 mL of 0.068 M potassium phosphate buffer (pH-6.8). The reaction began after 40 μ L of 10 mM H₂O₂ was added to the tube, and the optical density was measured at 240 nm. The change in absorbance at 240 nm in the linear phase of the slope ($\Delta 240 \text{ min}^{-1} \text{ mg}^{-1} \text{ protein}$) was used to measure catalase activity.

H₂O₂ content estimation

Alexieva *et al.* method⁴⁵ was used to determine the hydrogen peroxide concentration in plant leaves after reacting with KI, H₂O₂ was detected spectrophotometrically. 0.5 mL 0.1 percent trichloroacetic acid (TCA) leaf extract supernatant, 0.5 mL 100 mM K-phosphate buffer, and 2 mL reagent (1 M KI *w/v* in fresh double-distilled water) made up the reaction mixture. In the absence of leaf extract, the blank probe comprised of 0.1% TCA. The reaction was developed in the dark for 1 h, and the absorbance was measured at 390 nm. A standard curve with known H₂O₂ values was used to calculate the amount of hydrogen peroxide.

Malondialdehyde (MDA) content estimation

The amount of malondialdehyde (MDA), a result of unsaturated fatty acid peroxidation, can be used to quantify lipid peroxidation. The MDA concentration (mg g^{-1} fresh weight) was calculated using Li *et al.* technique⁴⁶. A total of 0.5 g of fresh leaves was homogenised in 10 mL of 10% trichloroacetic acid (TCA) and centrifuged at 12000 g for 10 min. An aliquot of 2 mL of the supernatant was then added to two millilitres of 0.6 percent thiobarbituric acid in 10% TCA. The combination was cooked for 30 min in boiling water before being immediately chilled in an ice bath. After centrifugation at 10000g for 10 min, the absorbance of the supernatant was measured at 450, 532, and 600 nm. The MDA concentration, which was expressed as $\text{nmol g}^{-1} \text{ DW}$, was estimated from the formula: $C (\text{nmol mL}^{-1}) = 6.45 (A_{532} - A_{600}) - 0.56(A_{450})$.

Protein assay

The protein content in the extract was estimated by the Bradford method using bovine serum albumin as standard.

Primer designing

For primer synthesis, the Primer3 Plus software (www.bioinformatics.nl/primer3plus) was utilised. The software was loaded with nucleotide sequences, qRT-PCR was selected as the reaction type, and a product size of 150–250 bp was chosen. The forward and reverse primers with the fewest variances in several parameters such as GC content, melting temperature, and self-complementarity were chosen, and specificity was validated using the NCBI nucleotide blast. The primers were designed in India by GCC Biotech Pvt. Ltd. For the experiment, the stock primers were diluted in a 1:10 ratio and maintained at -20°C . A total of seven pathogenicity-related primers were employed (Table 1⁴⁷⁻⁴⁹). For the amplification of *Rhizoctonia solani* cDNA, the housekeeping gene GADPH was included⁴⁷.

Gene expression studies

RNA isolation

Plants were watered two times a day, and inoculation was done in the evening. At 24, 48, 72, and 96 h after inoculation, the inoculated plants were harvested. The plants were labelled and covered in aluminium foil before being preserved at -80°C . The total RNA from infected rice sheaths was obtained using the Trizol technique (Invitrogen). All laboratory materials such as eppendorf tubes, spatulas, microcentrifuge tubes, pestle, and mortar were sterilised with DEPC (Diethyl pyro carbonate) treatment @ 0.1 percent for the inactivation of RNase enzymes. These materials were autoclaved for 15 min at 121°C and 15psi. Infected Pusa Basmati 1121 were homogenised with liquid N₂ in a pre-cooled mortar and pestle. To thoroughly dissociate the nucleoprotein complex, the fine powder (approximately 200 mg) was incubated in a 1.5 mL eppendorf tube with 1 mL Trizol reagent for 5 min at room temperature. For phase separation, the tubes were filled with chloroform (0.2 mL per 1 mL Trizol). The tubes were vigorously shaken by hand for 15 sec, allowed to cool for 2 to 3 min before being centrifuged at 12000 g for 15 min at 4°C . The aqueous phase was transferred to fresh tubes, pre-chilled with 0.5 mL isopropanol, and then centrifuged at 12000 g for 10 min at 4°C . By removing the supernatant solution from the tubes, the RNA pellet was maintained. The pellets were centrifuged at 7500 g for 5 min in 4°C after being rinsed with 1 mL of 75 percent ethanol. After removing the wash, the pellets were dried in air for 10 min. The pellets were dissolved in 30 μ L of DEPC-treated water and kept at

Table 1 —*Rhizoctonia solani* primers used for qRT-PCR analysis⁴⁷⁻⁴⁹

S. No.	Genes name	Primer name	Primer sequence
1	Inorganic phosphate transporter (IPT)	IPT-F	TCTTGACCACGGGGAGATAC
		IPT-R	GGATGTCCAGAGCAAACCAT
2	Bromodomain containing protein (BrD)	BrD-F	ATGCGCTCATTGAGGATTC
		BrD-R	CGGTAATCTACGCCCACT
3	Aldehyde dehydrogenase (AldD)	AldD-F	AATGGGCAGACTTGGATGAG
		AldD-R	GGAGGGTTATCGACAAACGA
4	AMP bindingdomain (AMP)	AMP-F	CAAGAAGGAGCTCGAAGTGG
		AMP-R	GGCGGCCACTTTTATTGTTA
		HmPr-F	GCTTGAACAAGACGAGGAG
5	Hemeperoxidase (HmPr)	HmPr-R	TTGAGAATAGGGCGAGGAGA
		LisH-F	GAGCTTACCTTGCACCTTG
6	Lissencephaly1 homolog (LisH)	LisH-R	ATGTAGATGTGCGGTGTGGA
		NIC-F	TGCGTTCTAACAGAGGGCGTA
7	Nuclear poreprotein, Nic96 (NIC)	NIC-R	ATTCTGGACTCAACCAAGC
		GAPDH-F	CAAAGTGACCAGCGACTTGA
8	Glyceraldehyde-3-Phosphate Dehydrogenase (GAPDH)	GAPDH-R	GCCTTGCTTGTGTCAGTGAAG

–80°C. The RNA was quantified using a Nano Drop-2000 Spectrophotometer (Thermo Fisher Scientific, USA).

cDNA synthesis

The Verso cDNA synthesis kit was used to transform the RNA template into cDNA according to the manufacturer guidelines (Verso cDNA, Thermo scientific, USA). OligodT primers were used to create first strand cDNA. In the C1000TM thermal cycler, reverse transcription settings consist of one cycle at 42°C for 30 min and one cycle of inactivation at 95°C for 2 min (Bio rad, USA). The cDNA was kept at a temperature of –20°C.

qRT-PCR analysis

The qRT- PCR samples were set up in triplicates, with a non-template control. For the reactions, 96-well qRT-PCR plates (Axygen, Inc., California) were used. Real-time PCR was performed using the Quant Studio 12K Flex real-time PCR machine (Applied Biosystems by Life Technologies, USA). The target gene amplification was monitored using DyNAmo flash SYBR (2X) green mix dye (Thermo Scientific, USA). To ensure the fluorescent dye's efficacy, the SYBR green mix dye was utilised in a dark environment. The total reaction volume was 20 µL, with 10 µL of SYBR, 1 µL of forward and reverse primers, 1 µL of template cDNA, and 7 µL of nuclease-free water. In qRT-PCR setup, Initial denaturation at 95°C for 10 min is followed by 40 cycles of denaturation at 95°C for 15 sec,

annealing at 51 to 54°C for 30 sec (different for each primer pair), extension at 72°C for 1 min, then default dissociation curve analysis to assess amplicon specificity. The target gene expression was quantified using the $2^{-\Delta\Delta C_t}$ technique using the reference gene GAPDH as a control⁵⁰. The pathogenicity gene average Ct values were subtracted from the average Ct values for the GAPDH gene. ΔC_t values are determined by subtracting values. After then, $\Delta\Delta C_t$ values for each treatment were determined by subtracting their individual ΔC_t values from the ΔC_t value of the negative control (calibrator). $\Delta\Delta C_t$ was calculated using the formula below:

$$\Delta\Delta C_t = (C_{t \text{ target}} - C_{t \text{ reference}})_{\text{Time } x} - (C_{t \text{ target}} - C_{t \text{ reference}})_{\text{Time } 0}$$

where, Time x = any time point and Time 0 = 1x expression of the target gene normalized to reference gene.

Finally, using the formula $2^{-\Delta\Delta C_t}$, the fold change expression data was derived. Standard deviation calculated from three technical replicates on each of three independent biological studies is represented by error bars.

Statistical analysis

Statistical analysis were done for all replicated samples by using OPSTAT software. Replications were maintained in a completely randomized design and all experiments were repeated twice and critical difference was calculated. For biochemical and gene

Table 2 — Phenotypic evaluation of rice genotypes treated with biocontrol agents

S.no.	Treatments	Lesion height (cm)	Plant height (cm)	Relative lesion height (%)	Number of tillers	Yield (g/m ²)
1	<i>Chaetomium globosum</i>	60.62	89.87	68.26	37.5	497
2	<i>Aspergillus niger</i>	63.87	84	75.75	26.25	458
3	<i>Talaromyces flavus</i>	63.12	76.87	82.45	31.62	464
4	<i>Pseudomonas fluorescens</i>	67.5	88.75	77.89	23.12	436
5	Control (R-359)	84.25	96	87.78	24.62	404
6	Critical difference	6.66	5.65	9.05	4.92	6.14

expression studies data are expressed as mean of 3 independent biological and 3 technical replications with \pm standard deviations.

Results

Evaluation of rice genotypes for sheath blight disease severity:

The disease severity of the *Rhizoctonia solani* virulent isolate (R-359) in Pusa Basmati 1121 was assessed using Relative Lesion Height (RLH %), plant height, lesion length, number of tillers, and grain yield. Different biocontrol agents, including as *Talaromyces flavus*, *Chaetomium globosum*, *Pseudomonas fluorescens*, and *Aspergillus niger*, showed significant differences in Pusa Basmati 1121 (Table 2). Inoculated control had the highest relative lesion height (87.78 cm), while treatment *Chaetomium globosum* had the lowest relative lesion height (68.26 cm). Treatment *Chaetomium globosum* had the highest average number of tillers (37.5), whereas, treatment *Pseudomonas fluorescens* had the lowest average number of tillers (23.12). The highest yield was found in *Chaetomium globosum* (497 g/m²), whereas the lowest yield was observed (436 g/m²) in plants treated with *Pseudomonas fluorescens*.

Biochemical analysis

N-acetylglucosamine polymer chitin is hydrolyzed by the chitinase enzyme. Chitinase enzyme lowers 3,5-dinitrosalicylic acid and forms a yellow colour product in the presence of N-acetylglucosamine polymer chitin⁴¹. Chitinase enzyme activity was measured in *Chaetomium globosum* treated plants at 48 hpi and *Pseudomonas fluorescens* treated plants at 72 hpi, with maximum enzyme activity (4.53 mol chitin min⁻¹ mg protein⁻¹) and minimum enzyme activity (0.030 mol chitin min⁻¹ mg protein⁻¹), respectively, (Fig 1A).

Laminarin is employed as a substrate for the glucanase enzyme in the β -1,3-glucanase enzyme assay, which yields reducing sugars. Dinitrosalicylic acid becomes a dark brown product when it combines

with the generated reducing sugar⁴². At 48 hpi, *Talaromyces flavus* had the highest enzyme activity (0.19 μ mol laminarin min⁻¹ mg protein⁻¹). At 24 hpi, plants treated with *Pseudomonas fluorescens* had the lowest enzyme activity (0.04 μ mol laminarin min⁻¹ mg protein⁻¹) (Fig 1B).

The production of cinnamic acid at 290 nm was used to test the phenylalanine lyase (PAL) assay. L-phenylalanine is converted to cinnamic acid by PAL. At 48 hpi, *Chaetomium globosum* treated plants had higher constitutive PAL enzyme activity (0.70 μ mol L-phenylalanine min⁻¹ mg protein⁻¹). At 72 hpi, plants treated with *Pseudomonas fluorescens* had the lowest enzyme activity (0.15 μ mol L-phenylalanine min⁻¹ mg protein⁻¹) (Fig 1C).

According to Bailly *et al.*⁴⁴, catalase enzyme assays were performed. The catalase enzyme uses H₂O₂ as a substrate. At 96 hpi, *Chaetomium globosum* treated plants had a greater catalase enzyme activity (60.16 μ mol H₂O₂ min⁻¹ mg protein⁻¹). At 72 hpi, plants treated with *Pseudomonas fluorescens* had the lowest enzyme activity (12.72 μ mol H₂O₂ min⁻¹ mg protein⁻¹) (Fig 1D).

H₂O₂ level was highest at 72 hpi (16.06 μ mol g⁻¹) in the infected control. At 48 hpi, the minimum H₂O₂ level in *Aspergillus niger* treatment (5.14 μ mol g⁻¹) (Fig 1E).

Malondialdehyde (MDA) levels were lowest at 24 hpi, in *Chaetomium globosum* treated plants (0.28 nmol mL⁻¹). At 96 hpi, *Talaromyces flavus* treated plants had a higher MDA concentration (0.94 nmol mL⁻¹) (Fig 1F).

Determination of total protein content

The Bradford method was used to determine the protein concentration of various materials (Table 3).

Relative expression analysis

In Pusa Basmati 1121, the expression pattern of pathogenicity-related genes of *Rhizoctonia solani* varied depending on the time and biocontrol agents used. The inorganic phosphate transporter (IPT) was marginally elevated at 96 hpi compared to 72 hpi




Fig 1 — Biochemical analysis of plant defence related enzymes in Pusa Basmati 1121 treated with biocontrol agents. (A)Chitinase activity, (B) β-1,3-glucanase activity, (C) Phenylalanine ammonia lyase activity, (D) Catalase activity, (E) Hydrogen peroxide (H₂O₂) content, and (F)Malondialdehyde (MDA) content

in all treatments, but relatively high expression at 96 hpi in the inoculated control (fold change 32.92) (Fig 2A).

In comparison to all other treatments, the expression pattern of Bromodomain containing protein (Brd) was highest in *Chaetomium globosum* at 96 hpi (fold change 39.36). At 72 hpi, all of the treatments had much lower expression, while at 96 hpi, they had significantly higher expression. In comparison to the other treatments, the inoculated control showed reduced expression (Fig 2B).

Inoculated control plants had significantly higher levels of heme peroxidase (HmPr) expression at 96 hpi (fold change 677.93) than other biocontrol treated plants. When compared to other treatments, the expression level of *Chaetomium globosum* was determined to be low at 96 hpi (fold change 8.33). In

addition, when compared to 96 hpi, there was very little expressiveness at 72 hpi (Fig 2C).

When comparing the inoculation control (*Rhizoctonia solani*) to the remainder of the treatment, the gene AMP binding domain (AMP) expression was found to be highest at 96 hpi (fold change 22.28) in the inoculated control (*Rhizoctonia solani*). When compared to expression at 72 hpi, all treatments had higher expression at 96 hpi (Fig 2D).

In comparison to other treatments, the expression of the nuclear pore protein Nic96 (NIC) was highest in *Pseudomonas fluorescens* treated plants at 96 hpi (fold change 238.36). When compared to the inoculated control (fold change 108.23), *Talaromyces flavus* demonstrated considerable expression at 96 hpi (fold change 193.74). When compared to other treated plants, *Chaetomium globosum* showed the least




Fig 2 — *Rhizoctonia solani* gene expression study using qRT-PCR in PB-1121 treated with biocontrol agents at distinct hours of post-inoculation. (A) Inorganic phosphate transporter (IPT), (B) Bromodomain function (BrD), (C) Heme peroxidase (HmPr), (D) AMP binding domain (AMP), (E) Nuclear pore protein, Nic96 (NIC), (F) Lissencephaly-1 homolog (LisH), and (G) Aldehyde dehydrogenase (AldD)

expression at both time intervals, at 72 hpi (fold change 22.70) and 96 hpi (fold change 24.36). At both time intervals, the inoculated control showed good levels of expression and were upregulated (Fig 2E).

At 96 h after inoculation, the Lissencephaly-1 homolog (LisH) was expressed at its maximum level in the infected control (fold change 652.12). In comparison to 96 hpi, the expression was higher at 72 hpi with the treatments *Aspergillus niger* (fold change 51.66), *Talaromyces flavus* (fold change 251.07) and *Pseudomonas fluorescens* (fold change 254.05) (Fig 2F).

The expression of aldehyde dehydrogenase (AldD) in the inoculated control was substantially identical to that of the other treatments. *Chaetomium globosum* (fold change 4.7), *Talaromyces flavus* (fold change

Treatments	Time Interval			
	24 hpi	48 hpi	72 hpi	96 hpi
Control(R-359)	80.741	50.74	87.40	108.14
<i>Chaetomium globosum</i>	25.556	85.92	170.37	85.55
<i>Aspergillus niger</i>	37.03	40.37	62.22	138.88
<i>Talaromyces flavus</i>	52.22	30.00	91.11	173.33
<i>Pseudomonas fluorescens</i>	76.66	63.33	82.59	97.03
Critical difference	0.002	0.004	0.004	0.002

11.42), and *Pseudomonas fluorescens* elevated expression levels at 96 hpi (fold change 9.64) (Fig 2G).

The inoculated control had a larger upregulation of pathogenicity-related genes, such as IPT, HmPr, AMP, LisH, and AldD, than the other treatments. At 96 hpi, genes such as Brd, IPT, HmPr, AMP, and NIC were found to be upregulated compared to 72 hpi.

Discussion

Rhizoctonia solani Kühn is a soil-borne polyphagous pathogen that wreaks havoc on a wide range of vital staple crops. *Rhizoctonia solani* Kuhn produces rice sheath blight, which reduces yield dramatically by causing necrotic lesions on the leaves, sheaths, and grains.

Different biological controls were applied, which aided in reducing the disease. Due to the harmful impacts of pesticides and agrochemicals on ecosystems, biological controls have been demonstrated to be helpful in reducing plant diseases⁵¹. There are roughly 35 species of fungi and bacteria that have been utilised as biocontrol agents against various plant infections as an alternative to agrochemicals⁵².

Biological controls, through physiological and biochemical changes, aid disease tolerance. In comparison to the inoculation control, rice plants treated with biocontrol agents had reduced relative lesion height and higher grain yield.

Chitinase is involved not only in plant defence, but also in plant development regulation⁵³. Plants generate enzymes that can break down pathogen cell walls, such as β -1,3-glucanase and chitinase⁵⁴. These enzymes are major determinants of plant fungal disease resistance⁵⁵. Hydrolytic enzymes act on fungal germlings shortly after pathogen penetration, weakening them and preventing disease development, whereas the pathogen may penetrate and colonise the tissue in susceptible hosts before PR proteins are induced to the needed level. The activity of chitinase, measured in mole of chitin $\text{min}^{-1} \text{mg protein}^{-1}$, was shown to be lowest in the inoculated control at 48 hpi when compared to all other treatments. As a result, it is obvious that chitinase activity in plants treated with biocontrol agents was often higher than inoculation control plants. It has been shown that increasing chitinase activity in plants treated with biological controls reduced pathogen damage⁵⁶. When compared to other time intervals, the enzyme activity were shown to be greater at 48 hpi. This could be related to plants' faster metabolic rate and increasing cells⁵⁷.

Pathogenic attacks, wounded tissue, UV radiation, heavy metal exposure, low temperature, and low amounts of nitrogen, phosphate, or ions are all factors that influence PAL activity⁵⁸. The stress response PAL induction and activity can be used as a regulatory control. As a result, by tracking PAL activity, it is possible to forecast and support the defence response of various biological controls.

Furthermore, PAL enzymes have a role in the synthesis of phenolic components that aid in the repair of plant wounds. Plants can use increased PAL activity as a defence mechanism to defend themselves against pathogen infection⁵⁹⁻⁶². The inoculation control treatment had the lowest PAL enzyme activity, according to the results of the investigation. Infection treatment combined with biological controls, on the other hand, results in increased activity, which was highest with the treatment *Chaetomium globosum*, showing that wounding induced PAL enzyme activity. This finding confirmed that biological regulations have an effect on the expression of PAL, a defence system component.

Any derivative of oxygen molecules (O_2) that is deemed more reactive than O_2 itself is known to as a reactive oxygen species (ROS). Thus, ROS encompasses both free radicals like superoxide (O_2^-) and hydroxyl radicals (OH) as well as non-radicals such singlet oxygen ($^1\text{O}_2$) and H_2O_2 . Because of its relative stability, H_2O_2 has drawn a lot of attention as a signal molecule involved in the regulation of particular biological processes, including interactions between plants and pathogens. As opposed to all of the treatments investigated, the amount of H_2O_2 in the inoculation control increased slightly. As a result, all of the biological controls utilised to control the level of H_2O_2 had a very modest effect.

CAT is a tetramericheme-containing protein that is arguably one of the most well-known H_2O_2 breaking enzymes. While plants possess a variety of H_2O_2 -metabolizing proteins, catalases seem to be highly active enzymes that catalyse a dismutase reaction and do not require cellular reductants. Catalases differ from many other peroxide-metabolizing enzymes in that they have a high selectivity for H_2O_2 but a low activity against organic peroxides⁶³. In comparison to the other treatments, the treatment *Chaetomium globosum* had the highest catalase activity. As a result, it is obvious that *Chaetomium globosum* was determined to be the most effective biological control for metabolising peroxides such as H_2O_2 .

To assess membrane damage in rice seedlings, lipid peroxidation in terms of malondialdehyde (MDA) was measured. When compared to the inoculation control and all other treatments, the treatment *Chaetomium globosum* had the lowest MDA content. To summarise, *Chaetomium globosum* was the most effective biological control for preventing rice seedling injury and yield loss.

Rhizoctonia solani, which infects rice, was studied in terms of gene expression. Most of the genes were dramatically raised in expression after rice infection in the susceptible variety Pusa Basmati 1121, and their expression altered over time. At 96 hpi, relative expression analysis of pathogenicity related genes such as inorganic phosphate transporter (IPT), heme peroxidase (HmPr), and AMP binding domain (AMP), all of which are involved in pathogen virulence, showed that inoculated control (32.92 folds IPT, 677.93 folds HmPr, and 22.28 folds AMP) was significantly higher than all other treatments. Heme peroxidase (HmPr) is a protein that helps to deactivate hydrogen peroxide. It aids pathogenicity inside the pathogen and immunity in the host. An acyl-activating enzyme (AAE) consensus motif is the AMP binding domain (AMP)⁶⁴ also expression level of these genes was upregulated at 96 hpi.

Bromodomain (BrD) expression was observed to be higher (39.36 times) in the *Chaetomium globosum* treatment, and it was also elevated at 96 hpi in all treatments. Many chromatin-associated proteins including nuclear histone acetyltransferases contain bromodomains. BrD-containing protein is an evolutionary conserved protein that plays a role in protein complex building via lysine acetylation and regulates gene transcription. This gene also plays a crucial role in cell proliferation. They predominantly interact with acetylated lysine⁶⁴.

Aldehyde dehydrogenase (AldD) is required for fungal pathogen development⁶⁵. When compared to the other treated plants, the expression pattern of the AldD gene was increased in inoculated control plants at 72 hpi (14.72) and 96 hpi (24.06).

Nuclear pore protein (NIC) is thought to play a role in pathogenesis since it is required for cell proliferation and gene regulation⁶⁶⁻⁶⁷. When 96 hpi is compared to 72 hpi, the expression of this gene is elevated in all treatments. At 96 hpi, *Pseudomonas fluorescens* displayed the most expression (238.36).

Through dynein and microtubule-dependent mechanisms, Lissencephaly-1 homolog (LisH) is crucial for mitotic spindle fibre synthesis and nuclear migration during cell division. Additionally, it engages in signal transduction via regulatory/adaptor systems, which are present in several eukaryotic proteins and are involved in the processing of pre-mRNA and the construction of the cytoskeleton⁶⁸. The majority of the treatments (*Aspergillus niger*, *Talaromyces flavus* and *Pseudomonas fluorescens*)

had higher expression at 72 hpi, however this was reduced at 96 hpi.

Conclusion

Biological management of Sheath blight disease has recently emerged as an appealing and promising alternative as use of chemicals are toxic for human health. Present investigation infers that the plant treated with biocontrol agents viz. *Chaetomium globosum*, *Aspergillus niger*, *Talaromyces flavus* and *Pseudomonas fluorescens* showed higher activity of plant defense related enzymes and also the expression of pathogenicity related genes were found to be less expressed in biocontrol treatments as compared to the inoculated control.

The findings of this study revealed that biological management of *Rhizoctonia solani* caused sheath blight disease in rice via induction of the defensive mechanism by plants provides insight into the plant-microbe interaction and a potential technique for plant protection and development in the future.

Acknowledgement

Authors are thankful to Head, Division of Plant Pathology, Joint Director (Research) and Director ICAR-IARI for providing research facilities.

Conflict of interest

All authors declare no conflict of interest.

References

- 1 Pathak HPMH, Samal P & Shahid M, Revitalizing rice-systems for enhancing productivity, profitability and climate resilience. Rice Research for Enhancing Productivity, Profitability and Climate Resilience. ICAR-National Rice Research Institute, Cuttack, Odisha, (2018) 1.
- 2 Turaidar V, Reddy M, Anantapur R, Krupa KN, Dalawai N, Deepak CA & Kumar KH, Rice sheath blight: major disease in rice. *Int J Curr Microbiol App Sci*, 7 (2018) 976.
- 3 Zhang SW, Yang Y & Li KT, Occurrence and control against rice sheath blight. *Biol Dis Sci*, 42 (2019) 87.
- 4 Molla KA, Karmakar S, Molla J, Bajaj P, Varshney RK, Datta SK & Datta K, Understanding sheath blight resistance in rice: the road behind and the road ahead. *Plant Biotechnol J*, 18 (2020) 895.
- 5 Savary S, Willocquet L, Elazegui FA, Castilla NP & Teng PS, Rice pest constraints in tropical Asia: quantification of yield losses due to rice pests in a range of production situations. *Plant Dis*, 84 (2020) 357.
- 6 Nagarajkumar M, Bhaskaran R & Velazhahan R, Involvement of secondary metabolites and extracellular lytic enzymes produced by *Pseudomonas fluorescens* in inhibition of *Rhizoctonia solani*, the rice sheath blight pathogen. *Microbiol Res*, 159 (2004) 73.
- 7 Wang LL, Liu LM, Hou YX, Li L & Huang SW, Pathotypic and genetic diversity in the population of *Rhizoctonia solani*

- AG 1-IA causing rice sheath blight in China. *Plant Pathol*, 64 (2015) 718.
- 8 Boukaew S & Prasertsan P, Suppression of rice sheath blight disease using a heat stable culture filtrate from *Streptomyces philanthi* RM-1-138. *Crop Prot*, 61 (2014) 1.
 - 9 Siddhartha, Verma A, Bashyal BM, Gogoi R & Kumar R, New nano-fungicides for the management of sheath blight disease (*Rhizoctonia solani*) in rice. *Int J Pest Manag*, 68 (2022) 217.
 - 10 Li D, Li S, Wei S & Sun W, Strategies to manage rice sheath blight: Lessons from interactions between rice and *Rhizoctonia solani*. *Rice*, 14 (2021) 1.
 - 11 Lee FN & Rush MC, Rice sheath blight: a major rice disease. *Plant Dis*, 67 (1983) 829.
 - 12 Pan XB, Rush MC, Sha XY, Xie QJ, Liscombe SD, Stenina SR & Orad JH, Major gene non-allelic sheath blight resistance from the rice cultivars Jasmine 85 and Teqing. *Crop Sci*, 39 (1999) 338.
 - 13 Rush MC & Lee FN, Sheath blight. In: *Compendium of rice diseases*, (Ed. by RK Webster & PS Gunnell; American Phytopathological Society, St. Paul, MN), (1992) 22.
 - 14 Zou JH, Pan XB, Chen ZX, Xu JY, Lu JF, Zhai WX & Zhu LH, Mapping quantitative trait loci controlling sheath blight resistance in two rice cultivars (*Oryza sativa* L.). *Theory Appl Genet*, (2000) 569.
 - 15 Mohammadi S & Ghanbari L, *In vitro* Antagonistic Mechanisms of *Trichoderma* spp. and *Talaromyces flavus* to Control *Gaeumannomyces graminis* var. *tritici* the Causal Agent of Wheat Take-all Disease. *Turk J Agric-Food Sci Technol*, 3 (2015) 629.
 - 16 Gholi-Nyakan M, Roostae A, Naraghi L & Heydari A, Investigating biological control of Fusarium wilt disease of cucumber caused by *Fusarium oxysporum* f.sp. *Cucumerinum* by *Talaromyces flavus* and *Trichoderma harzianum*. *Int J Agri Crop Sci*, 7 (2014) 154.
 - 17 Madi L, Katan T, Katan J & Henis Y, Biological control of *Sclerotium rolfsii* and *Verticillium dahliae* by *Talaromyces flavus* is mediated by different mechanisms. *Phytopathology*, 87 (1997) 1054.
 - 18 Naraghi L, Heydari A, Rezaee S, Razavi M & Afshari-Azad H, Biological control of *Verticillium* wilt of greenhouse cucumber by *Talaromyces flavus*. *Phytopathol Mediterr*, 49 (2010) 321.
 - 19 Marcia J, Biological control *Verticillium* wilt of eggplant in the field. *Plant Dis*, 66 (1982) 1166.
 - 20 Ojaghian MR, Potential of *Trichoderma* spp. and *Talaromyces flavus* for biological control of potato stem rot caused by *Sclerotinia sclerotiorum*. *Phytoparasitica*, 39(2011) 185.
 - 21 Catenulatum Gilman G, Biological control of *Sclerotinia* wilt of sunflower with *Talaromyces flavus* and *Coniothyrium minitans*. *Plant Dis*, 78 (1994) 231.
 - 22 Ashwini C, A review on *Chaetomium globosum* is versatile weapons for various plant pathogens. *J Pharmacogn Phytochem*, 8 (2019) 946.
 - 23 Zhang H & Yang Q, Expressed sequence tags-based identification of genes in the biocontrol agent *Chaetomium cupreum*. *Appl Microbiol Biotechnol*, 74 (2007) 650.
 - 24 Hoffland E, Hakulinen J & Van Pelt JA, Comparison of systemic resistance induced by avirulent and non-pathogenic *Pseudomonas species*. *Phytopathology*, 86 (1996) 757.
 - 25 Wei G, Kloepper JW & Tuzun, S, Induced systemic resistance to cucumber diseases and increased plant growth by plant growth promoting rhizobacteria under field condition. *Phytopathology*, 86 (1996) 221.
 - 26 Venkatasubbaiah P & Safeeulla KM, *Aspergillus niger* for biological control of *Rhizoctonia solani* on coffee seedlings. *Int J Pest Manag*, 30 (1984) 401.
 - 27 Niu DD, Liu HX, Jiang CH, Wang YP, Wang QY, Jin HL & Guo JH, The plant growth-promoting rhizobacterium *Bacillus cereus* AR156 induces systemic resistance in *Arabidopsis thaliana* by simultaneously activating salicylate- and jasmonate/ethylene-dependent signalling pathways. *Mol Plant-Microbe Interact*, 24 (2011) 533.
 - 28 Kurabachew H & Wydra K, Induction of systemic resistance and defense-related enzymes after elicitation of resistance by rhizobacteria and silicon application against *Ralstonia solanacearum* in tomato (*Solanum lycopersicum*). *Crop Prot*, 57 (2014) 1.
 - 29 Chandrasekaran M & Chun SC, Expression of PR-protein genes and induction of defense-related enzymes by *Bacillus subtilis* CBR05 in tomato (*Solanum lycopersicum*) plants challenged with *carotovora* subsp. *carotovora*. *Biosci Biotechnol Biochem*, 80 (2016) 2277.
 - 30 Jain S, Vaishnav A, Kumari S, Varma, A, Tuteja N & Choudhary DK, Chitinolytic *Bacillus*-mediated induction of jasmonic acid and defense-related proteins in soybean (*Glycine max* L. Merrill) plant against *Rhizoctonia solani* and *Fusarium oxysporum*. *J Plant Growth Regul*, 36 (2017) 200.
 - 31 Costanzo S, Jackson AK & Brooks SA, High-resolution mapping of Rsn1, a locus controlling sensitivity of rice to a necrosis-inducing phytotoxin from *Rhizoctonia solani* AG1-IA. *Theor Appl Genet*, 123 (2011) 33.
 - 32 Wang X, Jiang N, Liu J, Liu W & Wang GL, The role of effectors and host immunity in plant-necrotrophic fungal interactions. *Virulence*, 5 (2014) 722.
 - 33 Wei MM, Wang AJ, Liu Y, Ma L, Niu XY & Zheng AP, Identification of the novel effector RsIA_NP8 in *Rhizoctonia solani* AG1 IA that induces cell death and triggers defense responses in non-host plants. *Front Microbiol*, 11(2020) 1115.
 - 34 Abeles FB, Bosshart RP, Forrence LE & Habig WE, Preparation and purification of glucanase and chitinase from bean leaves. *Plant Physiol*, 47 (1971) 129.
 - 35 Wessels JGH & Sietsma JH, Fungal cell walls: a survey, in *Plant Carbohydrates II* (Springer, Berlin, Heidelberg) (1981) 352.
 - 36 Rawat K, Bashyal BM, Sharma S & Aggarwal R, *Talaromyces favus*: a potential biocontrol agent against the bakanae disease of rice, in *Ind phytopathol society 6th international conference on plant, pathogens and people' (challenges in plant pathology to benefit humankind IARI, New Delhi, India, 23-27 February) 2016*.
 - 37 Aggarwal R, Tewari AK, Srivastava KD & Singh DV, Role of antibiosis in the biological control of spot blotch (*Cochliobolus sativus*) of wheat by *Chaetomium globosum*. *Mycopathologia*, 157 (2004) 369.
 - 38 Kamil D & Sinha A, Production of endoglucanase and xylanase from indigenous strain of *Aspergillus niger* (ITCC no. 7790): optimization of cultural conditions. *Ind Phytopathol*, 64 (2011) 44.

- 39 Singh D, Yadav DK & Sinha S, Improvement of bio-efficacy of bacterial antagonists by using bleaching powder and resistant cultivars to control bacterial wilt of tomato. *The Horti Soc Ind*, 74 (2017) 552.
- 40 Bhaktavatsalam G, Satyanarayana K, Reddy APK & John VT, Evaluation for sheath blight resistance in rice. *Int Rice Res Newsl Int Rice Res Inst*, (1978).
- 41 Rojas-Avelizapa LI, Cruz-Camarillo R, Guerrero MI, Rodríguez-Vázquez R & Ibarra JE, Selection and characterization of a proteo-chitinolytic strain of *Bacillus thuringiensis*, able to grow in shrimp waste media. *World J Microbiol Biotechnol*, 15 (1999) 299.
- 42 Abeles FB & Forrence LE, Temporal and hormonal control of β -1, 3-glucanase in *Phaseolus vulgaris* L. *Plant Physiol*, 45 (1970) 395.
- 43 Mori T, Sakurai M & Sakuta M, Effects of conditioned medium on activities of PAL, CHS, DAHP synthase (DS-Co and DS-Mn) and anthocyanin production in suspension cultures of *Fragaria ananassa*. *Plant Sci*, 160 (2001) 355.
- 44 Bailly C, Leymarie J, Lehner A, Rousseau S, Côme D & Corbineau F, Catalase activity and expression in developing sunflower seeds as related to drying. *J Exp Bot*, 55 (2004) 475.
- 45 Alexieva V, Sergiev I, Mapelli S & Karanov E, The effect of drought and ultraviolet radiation on growth and stress markers in pea and wheat. *Plant Cell Environ*, 24 (2001) 1337.
- 46 Li X, Yang Y, Sun X, Lin H, Chen J, Ren J & Yang Y, Comparative physiological and proteomic analyses of poplar (*Populus yunnanensis*) plantlets exposed to high temperature and drought. *PLoS One*, 9 (2014) 107605.
- 47 Prashantha ST, Bashyal BM, Krishnan SG, Dubey H, Solanke AU, Prakash G & Aggarwal R, Identification and expression analysis of pathogenicity-related genes of *Rhizoctonia solani* anastomosis groups infecting rice. *Biotech*, 11 (2021) 12.
- 48 RiouxR, Manmathan H, Singh P, DelosReyes B, JiaY & Tavantzis S, Comparative analysis of putative pathogenesis-related gene expression in two *Rhizoctonia solani* pathosystems. *Curr Genet*, 57 (2011) 391.
- 49 Ghosh S, Gupta, SK & Jha G, Identification and functional analysis of AG1-IA specific genes of *Rhizoctonia solani*. *Curr Genet*, 60 (2014) 327.
- 50 Livak KJ & Schmittgen TD, Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods*, 25 (2001) 402.
- 51 Lockwood JL, Evolution of concepts associated with soilborne plant pathogens. *Ann Rev of Phytopathol*, 26 (1988) 93.
- 52 Cook RJ & Baker KF, The nature and practice of biological control of plant pathogens. *Am Phytopathological Soc*, 1983.
- 53 De Jong AJ, Cordewener J, Lo Schiavo F, Terzi M, Vandekerckhove J, Van Kammen A & De Vries SC, A carrot somatic embryo mutant is rescued by chitinase. *The Plant Cell*, 4 (1992) 425.
- 54 Santos IS, Machado OLT, Da Cunha M & Gomes VM, A chitinase from *Adenanthera pavonina* L. seeds: purification, characterization and immunolocalization. *Plant Sci*, 167 (2004) 1203.
- 55 Funnell DL, Lawrence CB, Pedersen JF & Schardl CL, Expression of the tobacco β -1, 3-glucanase gene, PR-2d, following induction of SAR with *Peronospora tabacina*. *Physiol Mol Plant Pathol*, 65 (2004) 285.
- 56 Broglie K, Chet I, Holliday M, Cressman R, Biddle P, Knowlton S, Mauvis CI & Broglie R, Transgenic plants with enhanced resistance to the fungal pathogen *Rhizoctonia solani*. *Science*, 254 (1991) 1194.
- 57 Graham MY, Weidner J, Wheeler K, Pelow ML & Graham TL, Induced expression of pathogenesis-related protein genes in soybean by wounding and the *Phytophthora sojae* cell wall glucan elicitor. *Physiol Mol Plant Pathol*, 63 (2003) 141.
- 58 Dixon RA & Paiva NL, Stress-induced phenylpropanoid metabolism. *Plant cell*, 7 (1995) 1085.
- 59 Chet I, Biotechnology in plant disease control. *Wiley Liss*, New York, 1993.
- 60 Seki H, Nagasugi Y, Ichinose Y, Shiraishi T & Yamada T, Changes in *in vivo* DNA-protein interactions in pea phenylalanine ammonialyase and chalcone synthase gene promoter induced by fungal signal molecules. *Plant Cell Physiol*, 40 (1999) 88.
- 61 Cui Y, Bell AA, Joost O & Magill C, Expression of potential defense response genes in cotton. *Physiol Mol Plant Pathol*, 56 (2000) 25.
- 62 Logemann E, Tavernaro A, Shulz W, Somssish I.E & Hahlbrock K, UV light selectively co induces supply pathway from primary metabolism and flavonoid secondary product formation in parsley. *Proc Natl Acad Sci U S A*, 97 (2000) 1903.
- 63 Mhamdi A, Queval G, Chaouch S, Vanderauwera S, Van Breusegem F & Noctor G, Catalase function in plants: a focus on Arabidopsis mutants as stress-mimic models. *J Exp Bot*, 61 (2010) 4197.
- 64 Zheng A, Lin R, Zhang D, Qin P, Xu L, Ai P, Sun Z & Li P, The evolution and pathogenic mechanisms of the rice sheath blight pathogen. *Nat Commun*, 4 (2013) 1424.
- 65 Wang X, Mann CJ, Bai Y, Ni L & Weiner H, Molecular cloning, characterization, and potential roles of cytosolic and mitochondrial aldehyde dehydrogenases in ethanol metabolism in *Saccharomyces cerevisiae*. *J Bacteriol*, 180 (1998) 822.
- 66 Grandi P, Doye V & Hurt EC, Purification of NSP1 reveals complex formation with 'GLFG' nucleoporins and a novel nuclear pore protein NIC96. *EMBO J*, 12 (1993) 3061.
- 67 Chu FH, Lee YR, Chou SJ, Chang TT & Shaw JF, Isolation and analysis of genes specifically expressed during basidiomatal development in *Antrodia cinnamomea* by subtractive PCR and cDNA microarray. *FEMS Microbiol Lett*, 280 (2008) 150.
- 68 Wibberg D, Andersson L, Rupp O, Goesmann A, Pühler A, Varrelmann M & Schlüter A, Draft genome sequence of the sugar beet pathogen *Rhizoctonia solani* AG2-2IIIB strainBBA69670. *J Biotechnol*, 222 (2016) 11.