



Blind docking of 4-Amino-7-Chloroquinoline analogs as potential dengue virus protease inhibitor using CB Dock a web server

Prasanna B Ranade^{1*}, Dinesh N Navale¹, Santosh W Zote², Dnyaneshwar K Kulal³ & Swapnil J Wagh⁴

¹Department of Chemistry, Vivekanand Education Society's College of Arts, Science and Commerce (Autonomous), Sindhi Society, Chembur, Mumbai 400 071, Maharashtra, India

²Department of Chemistry, PTVA's Sathaye College (Autonomous), Dixit Road, Vile Parle (East), Mumbai-400 057, Maharashtra, India

³Department of Chemistry, RamnarainRuia Autonomous College, L. N. Road, Matunga, Mumbai-400 019, Maharashtra, India

⁴Department of Chemistry, R.S.S. Prasarak Mandal's Nanasaheb Yashvantrao Narayanrao Chavan, Arts, Science & Commerce College Chalisgaon, Jalgaon-424 101, Maharashtra, India

Received 18 July 2022; revised 01 October 2022

Currently, there is no approved drug to combat dengue. Various quinoline derivatives are known for potential antimalarial, antiviral activities, *etc.* In the present work docking between 4-Amino-7-Chloroquinoline analogs was performed with dengue virus NS2B/NS3 protease using CB dock, a web server. Lys74, Ile165, Val147, Asn152, Asn167, Trp83 and Leu149 amino acid residues were found to be in contact with designed 4-Amino-7-Chloroquinoline analogs. Different modes of binding like hydrogen bonding, hydrophobic interactions, *etc.* with designed compounds improve potential anti-dengue characteristics *in silico*. ADME results are in acceptable range.

Keywords: 2FOM, ADME, Amino Acids, Drug design, Quinoline

Dengue is a viral disease caused due to the bite of *Aedes aegypti* type mosquito¹⁻². DEN 1, DEN 2, DEN 3 and DEN 4 are four serotypes of dengue. Dengue virus contains capsid, pre-membrane and envelope as structural proteins. Whereas, NS1, NS2A, NS2B, NS3, NS4A, NS4B, and NS5 are seven nonstructural proteins³. The dengue virus replication is due to cleavage between host cell protease and virus-encoded two component protease NS2B/NS3 proteins. The active site of dengue virus protease is located between His51, Asp75, and Ser135 residues³.

Quinoline derivatives show antiviral activity⁴. Moreover, the quinoline class also shows activity against the dengue virus⁵. Quinoline derivatives have potential anti-dengue characteristics against dengue *in silico* approach⁶. Substituted amino chloro quinoline derivatives were known to exhibit antiprotozoal⁷, anti-HIV⁸, anti-zika⁹, antimalarial¹⁰, and anticancer¹¹ activities. For the development of the new drug, docking is commonly used. It requires an understanding of the protein drug interactions. These interactions can be studied by using software like an autodock, glide, *etc.* Virtual screening of compounds can be easily done using docking¹²⁻¹⁴.

In the present work, the docking between new 4-Amino-7-Chloroquinoline derivatives and dengue virus protease is performed using CB Dock, a web-based server¹². We hope our study will be helpful in designing new entities against the dengue virus.

Experimental Section

Materials and Methods

Hardware

Molecular docking studies described herein were performed on Dell Inspiron 15R Laptop (Intel® Core™ i3-processor) running Windows 7 Home Basic Operating System.

Docking studies

Docking between ligands and protein were performed on CB Dock web server. (<https://cadd.labshare.cn/cb-dock2/php/index.php>)¹⁵.

Dengue virus NS2B/NS3 protease proteins were downloaded from www.rcsb.org website in *pdb* format. (PDB ID: 2FOM)¹⁶⁻¹⁷. Water molecules and other heteroatoms were removed automatically from protein when uploaded on CB Dock web server (Table 1).

Ligands Preparation

Ligands were drawn on Chemdraw software and saved in *.sdf* file format for docking using CB dock web server.

*Correspondence:

E-mail: prasannaranade@yahoo.com

Suppl. Data available on respective page of NOPR

Table 1 — Dock score of designed molecules

Sr No	Compound	R chain	Dock score	Hydrogen bonding, Hydrophilic interactions with following amino acids
1	Compound 1	H	-5.8	Lys74, Ile165, Val147, Asn152, Asn167, Trp83 and Leu149
2	Compound 2	Methyl	-6.9	Lys74, Ile165, Val147, Asn152, Asn167, Trp83 and Leu149
3	Compound 3	Ethyl	-7.3	Lys74, Ile165, Val147, Asn152, Asn167, Trp83 and Leu149
4	Compound 4	n-Propyl	-7.7	Lys74, Ile165, Val147, Asn152, Asn167, Trp83 and Leu149
5	Compound 5	n-Butyl	-8.0	Lys74, Ile165, Val147, Asn152, Asn167, Trp83 and Leu149
6	Compound 6	n-Pentyl	-8.3	Lys74, Ile165, Val147, Asn152, Asn167, Trp83 and Leu149
7	Compound 7	n-Hexyl	-8.9	Lys74, Ile165, Val147, Asn152, Asn167, Trp83 and Leu149
8	Compound 8	n-Octyl	-9.5	Lys74, Ile165, Val147, Asn152, Asn167, Trp83 and Leu149
9	Pinostrobin ³	-	-7.8	Lys74, Ile165, Val147, Asn152, Asn167, Trp83 and Leu149

Table 2 — Dock score of designed molecules

ADME Prediction							
Compound	H-bond acceptors	H-bond donors	Consensus Log P	GI absorption	BBB permeant	log Kp (cm/s)	Lipinski violations
1	1	1	2.1	High	Yes	-5.78	0
2	1	1	2.61	High	Yes	-5.05	0
3	1	1	2.94	High	Yes	-4.88	0
4	1	1	3.3	High	Yes	-4.59	0
5	1	1	3.63	High	Yes	-4.42	0
6	1	1	3.98	High	Yes	-4.13	0
7	1	1	4.32	High	Yes	-3.82	0
8	1	1	5.06	High	Yes	-3.22	0
9	4	1	2.66	High	Yes	-5.68	0

Swiss ADME program helped in computing the drug-likeness of designed compounds given in Table 1. Table 2 shows predictions of pharmacological properties. The pharmacokinetics of compounds that were analyzed using the descriptors and pharmaceutically relevant properties of ligands.

ADME Prediction

The swiss ADME program is used to predict physicochemical parameters and ADME parameters (<http://www.swissadme.ch/>) (Table 2).

Results and Discussion

Designed 4-Amino-7-Chloroquinoline derivatives were docked on dengue virus protease using CB dock web server. Previously reported anti-dengue compounds contains either hydrogen bonding or hydrophobic interactions with amino acids residues including Ile165, Lys74, Asn152, Leu149, Trp83, Ser135 and Val 147¹⁸⁻²⁴. Pinostrobin³ a flavanone derivative is known to exhibit anti-dengue activity with a docking score of -7.8 and it comes in contact with Lys74, Leu149, and Trp83 which are responsible for anti-dengue activity as per literature³.

Compound 1 to compound 9 were found to exhibit hydrogen bonding and/or hydrophilic interaction with Lys74, Leu149 and Trp83. It Indicates, the 4-Amino-7-Chloroquinoline class has potential for anti-dengue activity *in silico*. The docking score increases as the

carbon chain at the fourth position increases. The docking score is found to be comparable with Pinostrobin³.

Therefore, it can be concluded that the 4-Amino-7-Chloroquinoline class has a potential anti-dengue character.

The supporting images are provided in the supporting information file. The designed 4-Amino-7-Chloroquinoline compounds contact with amino acid residues is summarized in table.

Conclusion

It can be concluded that 4-Amino-7-Chloroquinoline analogs have potential anti-dengue activities *in-silico*. The interacting amino acid residues have a role in proteolytic cleavage in dengue replication, which is observed in designed 4-Amino-7-Chloroquinoline analogs indicating potential anti-dengue character. Blind docking helps in identifying potential target molecules for their activity against the dengue virus. CB dock program helps in predicting the potential anti-dengue compounds from dock score in less time.

Acknowledgement

Authors Prasanna Ranade and Dinesh Navale gratefully acknowledge the use of central instrumentation facilities at Vivekanand Education Society's College of Arts, Science and Commerce (Autonomous) funded by FIST-DST (Ministry of Science and Technology Govt. of India. and DBT STAR college scheme.

Conflict of interest

All authors declare no conflict of interest.

References

- Mira AMB, Christoph N, Veaceslav B & Christian DK, The Medicinal Chemistry of Dengue Virus. *J Med Chem*, 59 (2016) 5622.
- Jonathan C, Javier M, Soila SP, Michael SD & Rebeca RH, Mosquito Bite Delivery of Dengue Virus Enhances Immunogenicity and Pathogenesis in Humanized Mice. *J Virol*, 86 (2012) 7637.
- Rozana O, Tan SK, Norzulaani K, Rohana Y, Irene E, Newhouse JS, Newhouse MA & Noorsaadah AR, Docking of Noncompetitive Inhibitors into Dengue Virus Type 2 Protease: Understanding the Interactions with Allosteric Binding Sites. *J Chem Inf Model*, 48 (2008) 1582.
- Ramandeep K & Kapil K, Synthetic and medicinal perspective of quinolines as antiviral agents. *Eur J Med Chem*, 215 (2021) 113220.
- Carolina de la G, David ES, Hang TD, Mario Q, Oleg VL & Ricardo L, Antiviral Activity of Novel Quinoline Derivatives against Dengue Virus Serotype 2. *Molecules*, 23 (2018) 672.
- Chelsea NP & William NS, An *in silico* investigation of phytochemicals as antiviral agents against dengue fever. *Comb Chem High Throughput Screen*, 19 (2016) 516.
- Faist J, Hinteregger C, Seebacher W, Saf R, Mäser P, Kaiser M & Weis R, New derivatives of 7-chloroquinolin-4-amine with antiprotozoal activity. *Bioorg Med Chem*, 25 (2017) 941.
- Violetta C, Cristina P, Stefano M, Teresa P, Enrica F, Arianna C, Oriana T, Barbara G, Manlio P, Arnaldo F & Giorgio P, 6-Aminoquinolones as New Potential Anti-HIV Agents. *J Med Chem*, 43 (2000) 3799.
- Giselle BL, Ligia S da SP, Carlos RK, James LW, Caroline SF, Yasmine RV, Andressa M, José CN, Patrícia TB, Solange MS, Marcus VN & Thiago MLS, N-(2-(arylmethylimino) ethyl)-7-chloroquinolin-4-amine derivatives, synthesized by thermal and ultrasonic means, are endowed with anti-Zika virus activity. *Eur J Med Chem*, 127 (2017) 434.
- Murugan K, Panneerselvam C, Subramaniam J, Jayapal S, Manickam P, Rajapandian R, Murugan V, Jagannathan M, Syed S, Mathath R, Johan SPP, Stelia CM, Jonny ED, Al TA, Balamurugan C, Devakumar D, Shanmughavel P & Jiang-Shiou H, Synthesis of new series of quinoline derivatives with insecticidal effects on larval vectors of malaria and dengue diseases. *Sci Rep*, 12 (2022) 4765.
- Ferrer R, Gricela L, Gamboa N, Rodrigues J, Abramjuk C, Jung K, Lein M & Charris J, Synthesis of [(7-chloroquinolin-4-yl) amino] chalcones: Potential antimalarial and anticancer agents. *Sci Pharm*, 77 (2009) 725.
- Manisha C, Dolly S, Manas D & Karabi D, Molecular docking studies of natural and synthetic compounds against human secretory PLA2 in therapeutic intervention of inflammatory diseases and analysis of their pharmacokinetic properties. *Indian J Biochem Biophys*, 59, (2022)38.
- Jenny J & Kumar PB, Dihydroxyberberine from *Tinospora cordifolia*: *In silico* evidences for the mechanism of anti-inflammatory action through dual inhibition of Lipoxigenase and Cyclooxygenase. *Indian J Biochem Biophys*, 58 (2021) 244.
- Ganeshpurkar A & Saluja A, *In silico* interaction of hesperidin with some immunomodulatory targets: A docking analysis. *Indian J Biochem Biophys*, 56 (2019) 28.
- Yang L, Maximilian G, Wen-tao D, Mu-chun H, Zhi-Xiong X & Yang C, CB-Dock: a web server for cavity detection-guided protein-ligand blind docking. *Acta Pharmacologica Sinica*, 41 (2020) 138.
- Liangzhong L, Mei D, Amrita R, Jian K & Jianxing S, Curcumin Allosterically Inhibits the Dengue NS2B-NS3 Protease by Disrupting Its Active Conformation. *ACS Omega*, 40 (2020) 25677.
- Samina KY, Hasnah O, Mohammad SK, Basma M Abd R, Mohammed OE, Suriyati M, Othman S, Jualang AG & Thairarajan P, 4-Thiazolidinone coumarin derivatives as two-component NS2B/NS3 DENV favivirus serine protease inhibitors: synthesis, molecular docking, biological evaluation and structure- activity relationship studies. *Chem Central J*, 12 (2018) 69.
- Sriram RVU, Rajdip M, Annaram H, Sandip D, Achintya S, Uttam P, Nakul VR & Maiti C, Dabrafenib, idelalisib and nintedanib act as significant allosteric modulator for dengue NS3 protease. *PLoS One*, 10 (2021).
- Othman R, Othman R, Baharuddin A, Ramanan RN, Rahman Abd, N, Yusof R & Karsani S, Molecular Docking Studies of Selected Medicinal Drugs as Dengue Virus2 Protease Inhibitors. *Sains Malaysiana*, 46 (2017) 1865.
- Vijayakumar S, Ramesh V, Prabhu S, Nirajmohan S, Manogar. P, Natanamurugaraj G & Vathsala M, Virtual screening of phytochemicals that bind to dengue viral serotypes using molecular docking methods. *Int J Eng Res*, 9 (2018) 379.
- Tahir ul Q, Arooj M, Usman AA, Samia A, Tabeer F, Muhammad H, Syed SH, Waheed A & Sobia I, Computer Aided Screening of Phytochemicals from Garcinia against the Dengue NS2B/NS3 Protease. *Bioinformation*, 10(2014) 115.
- Sharoen Yu Ming Lim, Jin Yu Chieng & Yan P, Recent insights on anti-dengue virus (DENV) medicinal plants: review on *in vitro*, *in vivo* and *in silico* discoveries. *Pharmacol Pharm*, 14 (2021) 1.
- Sneha RC, Poonam AT, Rabindra KN & Sohan SC, Docking of Phytoconstituents of *Cynodon dactylon* on NS2B NS3 protease domain of Dengue virus. *Res J Pharm Tech*, 12 (2019) 5865.
- Sharmin S, Shafi M, Trina ET, Saad AS, Ahmad AA, Uzzal KA, Gobindo KP, Talha BE, Shahriar Z, Md. SU, Md. AS, Sultan A, Mohammed MG, Alruwali M, Ahmad JO, Nabilah RJ, Junghwan K & Bonglee KP, Phytochemical Compound Screening to Identify Novel Small Molecules against Dengue Virus: A Docking and Dynamics Study. *Molecules*, 27 (2022) 653.