Indian Journal of Biochemistry & Biophysics
Vol. 60, August 2023, pp. 601-610
DOI: 10.56042/ijbb.v60i8.4301

Ultrastructural variations of the sagittae in two *Anabantiformes* fishes: *Anabas testudineus* (Bloch, 1792) and *Trichogaster fasciata* (Bloch & Schneider, 1801)

Soumen Roy[1]*, Urmila Giri Roy[2], Narayan Ghorai[2], Samir Kumar Saha[2], Prabhakar Singh[3], Amit Mukhopadhyay[4] & Basudev Tripathy[5]

[1]Department of Zoology,City College, Kolkata-700 009, West Bengal, India

[2]Department of Zoology, West Bengal State University, Barasat, Kolkata-700 126, West Bengal, India

[3]Department of Anatomy, All India Institute of Medical Sciences, New Delhi-110 029, Delhi, India

[4]Mollusca Section, Zoological Survey of India, Kolkata-700 053, West Bengal, India

[5]Western Regional Centre, Zoological Survey of India, Pune-411044, Maharashtra, India

Received 15 February 2023; *revised 15 May 2023*

Supplementary Table

Table S1 — Relatedness matrix of the *Anabas testudineus*, *Trichogaster fasciata*, and five other fish species in the order *Anabantiformes* (*Trichogaster lalia*[1], *Channa limbata*[2], *Channa lucius*[2], *Channa micropeltes*[2], *and Channa striata*[2]) and a fish species of the order Perciformes (Perca fluviatilis; Yilmaz *et al*., 2014) on the sagitta morphostructural characteristics. [1]Banu and Sirajuddin, 2020; [2]Krainara and Wunnitikul, 2004

Correlation/ Relatedness matrix	*Anabas testudineus*	*Trichogaster fasciata*	*Trichogaster lalia*[1]	*Channa limbata*[2]	*Channa lucius*[2]	*Channa micropeltes*[2]	*Channa striata*[2]	*Perca fluviatilis*[2]
Anabas testudineus	0	4	4	10	10	10	9	8
Trichogaster fasciata		0	11	4	4	3	4	2
Trichogaster lalia[1]			0	4	4	3	4	2
Channa limbata[2]				0	11	9	9	8
Channa lucius[2]					0	9	9	8
Channa micropeltes[2]						0	9	8
Channa striata[2]							0	8
Perca fluviatilis[2]								0