

Supplementary Information

Syntheses and characterizations of 2-nitrobenzaldehyde amino acid Schiff base reductive amination compounds

Yu-Rou Jiang^a, Dan Su^a, Mu-Zhao Wang^b, Ai-Quan Jia^a & Qian-Feng Zhang^{a,*}

^aInstitute of Molecular Engineering and Applied Chemistry, Anhui University of Technology, Ma'anshan, Anhui 243002, P. R. China

^bDepartment of Chemistry, University of Warwick, Library Road, Coventry, CV4 7AL, UK

*E-mail: zhangqf@ahut.edu.cn

Sl. No.	Contents	Pg. No.
1	Fig. S1 — ¹ H NMR spectrum of aspartic acid	2
2	Fig. S2 — ¹ H NMR spectrum of glutamic acid	2
3	Fig. S3 — ¹ H NMR spectrum of leucine	3
4	Fig. S4 — ¹ H NMR spectrum of valine	3
5	Fig. S5 — FTIR spectrum of aspartic acid	4
6	Fig. S6 — FTIR spectrum of glutamic acid	4
7	Fig. S7 — FTIR spectrum of leucine	5
8	Fig. S8 — FTIR spectrum of valine	5




Fig. S1 — ^1H NMR spectrum of aspartic acid




Fig. S2 — ^1H NMR spectrum of glutamic acid




Fig. S3 — ¹H NMR spectrum of leucine




Fig. S4 — ¹H NMR spectrum of valine




Fig. S5 — FTIR spectrum of aspartic acid




Fig. S6 — FTIR spectrum of glutamic acid




Fig. S7 — FTIR spectrum of leucine




Fig. S8 — FTIR spectrum of valine