

## Aerobic Palladium(II) catalysed oxidative route for synthesis of Papaveraldine

Hari K. Kadam,<sup>a\*</sup> Tushar Anvekar,<sup>b</sup>


<sup>a</sup> School of Chemical Sciences, Goa University, Taleigao Plateau, Goa – 403206

<sup>b</sup> Department of Chemistry, St. Xavier's College, Mapusa, Goa – 403507


\*harikadam05@gmail.com, harikadam@unigoa.ac.in

### SUPPORTING INFORMATION

Commercial reagents were used without further purification. Solvents were distilled prior to use and dried if necessary, using standard procedures. Thin layer chromatography (TLC) was carried out on Silica gel 60 F254 aluminium plates purchased from Merck and were developed in iodine or UV chamber. All melting points and boiling points were recorded using Thiele's tube and are uncorrected. IR spectra were recorded on Shimadzu FT-IR spectrophotometer. <sup>1</sup>H NMR (400 MHz) and <sup>13</sup>C NMR (100 MHz) spectra were recorded on a Bruker AVANCE 400 MHz instrument and the multiplicities of carbon signals were obtained from DEPT experiment. LCMS were recorded on Agilent Technologies instrument.



HK-LM-08 CMR CDCL3



HK-LM-08 DEPT CDCL3

