

Supplementary Information

Copper sulphate immobilized on P(AN-NIPAM-MBAM) terpolymer as a highly efficient catalyst for the selective reduction of nitro-arenes

Anirudhdha G Kalola, Pratibha Prasad, Roshni D Hingrajiya & Manish P Patel*


Department of Chemistry, Sardar Patel University, Vallabh Vidyanagar 388 120, Gujarat, India

E-mail: patelmanish1069@yahoo.com, anirudhdha77patel@gmail.com

Received 15 September 2022; accepted (revised) 18 May 2023


Sr. no.	Content	Page no.
1	¹ H NMR & ¹³ C NMR of 2-chloroaniline	2
2	¹ H NMR & ¹³ C NMR of 4-chloroaniline	3
3	¹ H NMR & ¹³ C NMR of 4-bromoaniline	4
4	¹ H NMR & ¹³ C NMR of 4-iodoaniline	5
5	¹ H NMR & ¹³ C NMR of 4-aminophenol	6
6	¹ H NMR & ¹³ C NMR of 2,5-diethoxyaniline	7
7	¹ H NMR & ¹³ C NMR of 4-aminobenzoic acid	8
8	¹ H NMR & ¹³ C NMR of methyl 2-aminobenzoate	9
9	¹ H NMR & ¹³ C NMR of 4-aminobenzamide	10
10	¹ H NMR & ¹³ C NMR of N-(4-aminophenyl)acetamide	11
11	¹ H NMR & ¹³ C NMR of 4-aminobenzonitrile	12

¹H NMR of 2-chloroaniline





¹³C NMR of 2-chloroaniline

¹H NMR of 4-chloroaniline





^{13}C NMR of 4-chloroaniline

^1H NMR of 4-bromoaniline





^{13}C NMR of 4-bromoaniline

^1H NMR of 4-iodoaniline





^{13}C NMR of 4-iodoaniline

^1H NMR of 4-aminophenol





¹³C NMR of 4-aminophenol

¹H NMR of 2,5-diethoxyaniline





¹³C NMR of 2,5-diethoxyaniline

¹H NMR of 4-aminobenzoic acid





¹³C NMR of 4-aminobenzoic acid

¹H NMR of methyl 2-aminobenzoate





¹³C NMR of methyl 2-aminobenzoate
¹H NMR of 4-aminobenzamide





¹³C NMR of 4-aminobenzamide

¹H NMR of N-(4-aminophenyl)acetamide



¹³C NMR of N-(4-aminophenyl)acetamide

¹H NMR of 4-aminobenzonitrile



^{13}C NMR of 4-aminobenzonitrile

