

## Marine polysaccharides and potential enzymes for its degradation

Bhupendra Baghel<sup>1,#</sup>, Reeta Rani Singhania<sup>1,2,#,\*</sup>, Anil Kumar Patel<sup>1,2</sup>, Navneet Kurrey<sup>3</sup>, Chiu-Wen Chen<sup>1,2</sup> & Cheng-Di Dong<sup>1,2,\*</sup>

<sup>1</sup>Institute of Aquatic Science and Technology, National Kaohsiung University of Science and Technology, Kaohsiung City-81157, Taiwan


<sup>2</sup>Centre for Energy and Environmental Sustainability, Lucknow 226 029, Uttar Pradesh, India

<sup>3</sup>School of Engineering, Ajeenkya DY Patil University, DY Patil Knowledge City, Pune, Maharashtra, India

<sup>4</sup>Department of Marine Environmental Engineering, National Kaohsiung University of Science and Technology, Kaohsiung City, 81157 Taiwan

Received 31 January 2024; revised 21 March 2024

### Supplementary Figure



Suppl. Fig. S1 — Chemical structure of (A) Carrageenan; (B) Fucoidan; (C) Ulvan; (D) Chitosan; (E) Chitin; (F) Alginate; (G) Agarose; and (H) Porphyran