

Impact of binocular vision anomalies and stereopsis on individual performance: A comprehensive review of binocular vision disorders

Astha Mishra¹, Shankar Dass², Bunty Sharma³ & Ujjawal Sharma^{4*}

¹Chitkara School of Health Sciences, Chitkara University, Rajpura, Punjab 140401, India

²Optometrist, Health Department, Haryana Government, India

³Department of Biotechnology, Graphic Era (Deemed to be University), Dehradun, Uttarakhand, India

⁴Department of Human Genetics and Molecular Medicine, Central University of Punjab, Bhatinda 151001, India

Received 20 November 2025; revised 19 December 2025

Binocular vision disorders and anomalies in stereopsis primarily affect individuals' everyday functioning, academic performance, and work efficiency. Binocular vision is the coordination between the two eyes and is essential for depth perception, spatial awareness, and visual acuity. Strabismus, amblyopia, convergence insufficiency, and suppression can cause defective stereopsis, visual exhaustion, double vision, and poor reading speed. These conditions typically affect tasks that require eye coordination, such as driving, sports, and fine motor tasks. Patients with binocular vision disorders typically complain of headaches, asthenopia, and difficulty sustaining focus, which can result in decreased efficiency in school and work environments. In children, undiagnosed binocular vision abnormalities can be a cause of learning problems, misattributed as attention disorders or dyslexia. Occupations that require precise depth perception, such as aviation, surgery, and engineering, can be significantly impacted by stereopsis deficits, compromising overall performance and safety. Early diagnosis and treatment, such as vision therapy, corrective lenses, and surgery, can mitigate the negative consequences of these disorders. The current review examines the effects of binocular vision anomalies on daily performance and highlights the importance of early diagnosis and management techniques to improve individuals' quality of life. Educating educators, employers, and healthcare providers about these disorders is essential in creating an accommodating and supportive environment for individuals with these disorders.

Keywords: Binocular anomalies, Stereopsis, Binocular vision, convergence insufficiency, Accommodative anomalies

Introduction

Binocular single vision is the ability to see with both eyes at the same time. This is done by coordinating the use of both eyes. This makes it possible for two different but similar images to be seen as one image through the process of fusion^{1,2}. This system integrates motor alignment, sensory fusion, and cortical processing, allowing individuals to perceive their environment in three-dimensional space³.

A Stereoacuity is the smallest difference in retinal image that may be seen, which allows for the impression of fine depth differences. It is a key part of binocular vision. Stereoacuity enables individuals to make accurate spatial judgments essential for daily activities and professional responsibilities, such as reading, driving, object manipulation, sports, and careers that require advanced visuomotor coordination^{3,4}. Higher stereoacuity levels are the

cause of higher accuracy in visually guided tasks such as reading, driving, sports, and fine motor skills-related occupations such as surgery and engineering, early diagnosis and appropriate intervention, including vision therapy and correction of refractive error, play crucial roles in augmenting stereoacuity and its impact on occupational performance and activities of daily living⁴. Good stereopsis relies on several factors coming together: accurate ocular alignment, equal visual acuity between the eyes, healthy retinal correspondence, and intact binocular neural pathways. Any disturbance in these elements, usually from refractive errors, anisometropia, amblyopia, strabismus, or convergence insufficiency, can lower or abolish stereopsis. Clinically, normal stereoacuity would indicate intact binocular function and usually excludes significant ocular misalignment or amblyopia. On the other hand, even small deviations in eye posture or accommodation can degrade stereoacuity, which demonstrates the sensitivity of this clinical marker³⁻⁵.

*Correspondence:
E-mail: Ujjawal.sharma@cup.edu.in

Epidemiological studies estimate that about 3-5% of the population is stereoblind, meaning they cannot perceive depth due to binocular disparity. A larger percentage exhibits stereo anomalies, in which stereopsis is present but severely reduced. These conditions generally arise from disturbances in the early-life binocular visual system, such as untreated strabismus or amblyopia that occurred during critical developmental periods. If left untreated, these anomalies may persist throughout life and significantly impair the depth perception required for everyday activities⁶⁻⁸. The rapid advancement of digital technology in recent years has introduced new considerations related to binocular vision. Widespread exposure to three-dimensional (3D) media, including cinema, gaming, virtual simulations, and educational displays, has increased visual demands on the vergence and accommodation systems. Unlike conventional two-dimensional (2D) images, 3D displays require the eyes to converge and focus at different depths, a mismatch known as vergence accommodation conflict.

Excessive or prolonged exposure can induce 3D asthenopia, characterised by eye strain, headache, dizziness, blurred vision, and general visual fatigue.⁷⁻¹¹ Individuals with pre-existing binocular anomalies tend to experience such discomfort more intensely and more frequently. Understanding these responses is essential as 3D technologies become embedded in entertainment, medicine, military training, and industrial automation^{10,11}. Physiological studies have demonstrated that, in individuals with abnormal binocular function, even those with measurable stereopsis, reports exist of greater ocular fatigue, slower task completion, and reduced visual performance when working with stereoscopic stimuli. Poor stereopsis in some individuals sometimes brings about compensatory mechanisms, like better reliance on monocular cues or enhanced attention to the motion and texture of objects, thereby demonstrating the adaptability of the visual system¹². However, these compensations are often insufficient for tasks requiring fine depth discrimination, and affected individuals may remain unaware of their limitations due to the subtle nature of stereoscopic deficits.¹²⁻¹⁴ Importantly, because depth perception relies on multiple cues, both binocular (e.g., disparity, convergence) and monocular (e.g., motion parallax, perspective), individuals with mild binocular dysfunction may not recognise their deficit, resulting in delayed diagnosis and intervention.

Disorders such as convergence insufficiency, accommodative dysfunction, or intermittent strabismus may present with intermittent symptoms, complicating early detection¹⁴⁻¹⁶. Timely assessment of stereoacuity, therefore, plays a crucial role in both clinical decision-making and determining an individual's functional performance in visually demanding environments¹⁴.

The purpose of this review is to summarize the current evidence regarding the types of binocular vision disorders, their underlying mechanisms, and the impact on daily function, with an emphasis on the importance of early diagnosis and proper therapeutic or corrective measures.

Methodology

This narrative review compiles the evidence concerning binocular vision anomalies and their effects on functional performance. A selective search of pertinent peer-reviewed articles from prominent scientific databases, including PubMed, Google Scholar, Scopus, and Web of Science, guided the literature identification process. For this review, we focused on clinically relevant, well-designed studies about binocular vision problems, stereopsis, depth perception, and how they affect school, work, and everyday life. To get a maximum coverage of the topic being studied, both classical, modern and the most recent 2011 onwards works were included. A formal systematic search strategy and PRISMA-based reporting have not been conducted, as this is a narrative review.

Discussion

This review outlines the clear functional impact of binocular vision anomalies and reduced stereopsis on daily and task-specific performance. Although these often go unrecognized, they nevertheless show consistent effects on visual efficiency and depth perception. The discussion integrates key findings and their clinical implications.

Stereoacuity and binocular vision disorders

Abnormalities of motor alignment disrupt the normal correspondence between the two retinal images, while sensory abnormalities affect the neural integration necessary for processing depth. Impaired stereopsis, prolonged reaction times in depth-related tasks, and decreased accuracy in spatial judgments are common features of binocular vision disorders¹⁵. Even small losses in stereoacuity can significantly impact

visual efficiency, especially under conditions of high demands on rapid depth estimation or fine hand-eye coordination¹⁴. In more severe cases, stereoacuity may be completely absent. This occurs most frequently in conditions such as constant strabismus and severe amblyopia, where the brain suppresses input from the deviating or weakened eye to avoid diplopia. Prolonged suppression alters the developmental course of binocular pathways and, if left untreated during early childhood, may lead to permanent stereoblindness. Subjects with stereoblindness must always rely on monocular cues, which are much less accurate for near-depth tasks. They tend to move cautiously or more slowly in activities that demand in-depth judgment and commonly report problems with spatial localization, participating in athletics, and actions that require combined motor responses¹⁷.

Functional consequences extend beyond real-world contexts to interactions with digital environments. Individuals with binocular anomalies commonly report increased visual discomfort when viewing stereoscopic displays or performing virtual reality tasks that require accurate vergence responses¹²⁻¹⁶. There is evidence to suggest that degraded stereoacuity is strongly associated with heightened visual fatigue, fixation instability, and increased accommodative effort when viewing under such conditions; thus, assessing binocular status is important in contexts involving three-dimensional visual stimuli¹⁶. The loss of binocular vision, therefore, considerably impairs stereoacuity, the extent of which depends on the causing factor, the severity, and the duration of the condition. This is essential to understand the fine interactions among them for the exact diagnosis, detailed functional evaluation, and planning appropriate management to restore or optimize the stereoscopic function¹⁷⁻¹⁹(Table 1).

Binocular vision anomalies' impact on an individual's performance

Binocular vision anomalies can impact academic, occupational, and daily functioning in a manner that is

frequently subtle yet functionally significant. Many people, especially children, may go undiagnosed due to the similarity of their symptoms with other non-visual conditions, such as inattention or learning delays¹⁹. Children with poor binocular coordination need to exert more mental effort to maintain single vision, resulting in less cognitive reserve available for comprehension and sustained attention. Such an increased visual load may appear as reduced reading fluency, avoidance of near tasks, or inconsistent academic engagement, which could therefore be mistakenly attributed to behavioural or learning disorders²⁰.

The functional consequences in adults depend to a great degree on the precision demands of their visual environment. Those jobs requiring fine depth discrimination and visually guided motor skills are particularly sensitive to small stereo deficits. Even when individuals adapt to using monocular cues, such compensations are slower and less accurate, resulting in increased error rates, fatigue, and performance variability. Reduced binocular stability may also elevate occupational stress since tasks that should feel automatic require deliberate visual effort²⁰⁻²².

These functional limitations stem not only from the absence of stereopsis but also from the inefficiency of substituting less reliable monocular cues for rapid depth judgments²³⁻²⁵. Overall, binocular vision anomalies affect more than individual tasks—they reduce overall visual efficiency and disrupt cognitive-motor coordination²². The ability to detect this early and treat appropriately is extremely important in preventing long-term functional limitations and enhancing quality of life²¹⁻²³.

Academic challenges

Students with binocular vision disorder tend to experience difficulties in reading, writing, and other visually intensive schoolwork. Reading more slowly, loss of place too often, low comprehension, and more fatigue can result from trouble tracking words on a page. Students will, therefore, begin to dislike reading, which can impact the learning process in general.

Table 1 — Binocular Vision Anomalies and Their Effects

Disorder	Description	Impact on Performance
Strabismus	Misalignment of the eyes, leading to impaired fusion.	Difficulty in depth perception, visual discomfort.
Amblyopia (Lazy Eye)	Reduced vision in one eye due to abnormal development.	Decreased stereoacuity, reading difficulties.
Convergence Insufficiency	Difficulty in maintaining focus at near distances.	Visual fatigue, headaches, reading issues.
Suppression	Brain ignores one eye's image to prevent double vision.	Poor depth perception, reduced visual awareness.
Aniseikonia	Difference in perceived image size between the eyes.	Discomfort, difficulty in fusing images.

Table 2 — Binocular Vision Anomalies Treatment Methods

Treatment Method	Description	Conditions Treated
Vision Therapy	Orthoptic exercises to improve eye coordination, convergence, and depth perception.	Convergence Insufficiency, Accommodative Dysfunction, Strabismus
Corrective Lenses & Prisms	Glasses or contact lenses correct refractive errors, and prisms assist in binocular vision alignment.	Myopia, Hyperopia, Astigmatism, Strabismus, Aniseikonia
Surgical Interventions	Surgical procedures to align eye muscles and restore binocular function.	Severe Strabismus, Ocular Misalignment
Digital Vision Therapy	Virtual reality-based therapy for enhancing stereopsis and binocular coordination.	Amblyopia, Convergence Insufficiency, Binocular Dysfunction
Ergonomic Accommodations	Improved lighting, screen positioning, and scheduled breaks to reduce eye strain.	Eye Strain, Accommodative Dysfunction, Computer Vision Syndrome

Studies show that some children with undiagnosed binocular vision problems may experience difficulty focusing and paying attention, which can be misinterpreted as learning disorders or behavioural problems. These difficulties can lead to frustration, poor academic performance, and low self-confidence in the classroom²⁶.

Occupational implications

Vocational activities that require large amounts of visual-motor ability, such as flying pilots, surgical doctors, engineering experts, and sportspeople, utilize the best binocular vision for accuracy and reliability²⁷⁻²⁹. Binocular vision defects compromising depth perception contribute to mistakes, less efficiency, and even jeopardy to safety in many cases. For landing and moving aircraft safely in aviation fields, pilots benefit from good judgment of depth; surgeons in need of unerringly sure hand-eye movements during intricate medical operations. Similarly, construction, manufacturing, and machinery workers must accurately estimate distances to ensure safety and maintain efficiency. Stereopsis problems in these professions can heighten the chances of accidents and affect overall performance³⁰.

Social and Psychological Impact

In addition to academic and work-related challenges, binocular vision disorders can influence a person's self-esteem and social relationships. Disorders like strabismus, in which the eyes are not aligned, can result in apparent visual differences that lead to social stigma, embarrassment, and lower confidence. Children and adults with apparent eye alignment problems can be teased, bullied, or self-conscious in social situations. In addition, chronic visual distress, headaches, and fatigue may lead to stress and anxiety, prompting individuals to avoid

tasks necessitating prolonged visual attention, such as reading, sports, or social interaction. These psychological impacts underscore the importance of early diagnosis and treatment in enhancing functional vision and overall quality of life.

Diagnosis and Treatment

Early detection and management of binocular vision disorders are paramount in reducing their effect on performance^{31,32}. Diagnostic instruments, such as the Titmus test, Randot stereo test, and Worth 4-dot test, measure stereoacuity and binocular function Table 2. Other diagnostic tests include.

Cover Test

To determine ocular misalignment, it is comprised of two primary procedures. The Unilateral Cover Test, which identifies tropias (manifest deviations) by seeing if the uncovered eye will move to reacquire fixation, and the Alternating Cover Test, which detects phorias (latent deviations) by quickly alternating an occluder between the eyes and observing any movement. Detecting these misalignments is crucial because they can cause eye strain, double vision, and difficulties with depth perception, which affect reading, coordination, and daily visual tasks³³.

Vergence Testing

Measures the convergence and divergence ability of the eyes. Vergence Testing assesses the capacity of the eyes to converge (drive inward) or diverge (move outward) in order to produce clear single vision. Convergence Testing establishes whether the eyes converge efficiently to a near point when focusing on a near object. Drifting away of one eye or double vision when the patient is observed may suggest convergence insufficiency (CI), a condition associated with headaches, visual discomfort, and reading

problems. Divergence Testing, however, evaluates the ability of the eyes to move outward when viewing a far object. Inability to do so may lead to divergence insufficiency, resulting in occasional double vision and difficulty sustaining focus on distant objects. Vergence dysfunctions tend to make reading, driving, and long periods of screen time difficult^{34,35}.

Accommodation Tests

Determine the focusing capability of the eyes at varying distances. The Accommodative Facility Test measures the rate of change of accommodation between far and near points, using plus and minus lenses to assess accommodative facility, which is often symptomatic with fatigue. The final test is the Accommodative Response Test, which assesses the functionality of the eyes by retaining focus on an object at a constant distance to identify conditions such as accommodative spasm or insufficiency that can hinder reading efficiency and comfort³⁶.

Treatment Options

Vision Therapy:

Orthoptic exercises to enhance eye coordination, convergence, and depth perception. Vision Therapy is a non-surgical method that includes orthoptic exercises to develop eye coordination, convergence, and depth perception. Vision Therapy comes in handy when dealing with eye conditions like convergence insufficiency and accommodative dysfunction, in which patients develop their binocular vision through proper exercises conducted under professional guidance³⁷.

Corrective Lenses and Prisms

Refractive errors can be corrected with prescription glasses or contact lenses, and prisms can assist in binocular vision alignment. These are frequently applied to correct refractive errors and facilitate binocular vision alignment. Contact lenses or prescription glasses correct visual defects like myopia, hyperopia, and astigmatism, while prisms realign the eyes and reduce tension in patients with binocular vision defects. The treatments can significantly enhance comfort and visual acuity, ultimately improving performance in everyday tasks³⁸.

Surgical interventions

In extreme cases of strabismus, surgery may be required to align the eyes and restore binocular vision.

In severe misalignment, Surgical Interventions may be necessary, particularly for strabismus. Surgery is aimed at realigning the eye muscles and returning binocular function to normal. While not necessarily the first course of treatment, surgical intervention can be a wonderful help to those with persistent ocular deviations that are not responsive to other therapies³⁹.

Digital vision therapy

New technologies, including virtual reality-based therapy, offer novel solutions for enhancing stereopsis and binocular coordination. Virtual reality therapy, for example, enables patients to participate in interactive exercises designed to stimulate both eyes and enhance depth perception. These digital options offer interactive, adaptive, and efficient rehabilitation techniques that can be tailored to specific requirements. With the progress of technology. Digital Vision Therapy has become a promising option for improving stereopsis and binocular coordination⁴⁰.

Ergonomic accommodations

Accommodations like enhanced illumination, monitor positioning, and planned breaks can allow individuals with eye strain to successfully control their situation^{39,41}.

Anomalies of binocular vision pose significant difficulties in several areas of life, such as education, work, and psychological stability^{42,43}. Comparing students with normal binocular vision and individuals with anomalies, the academic achievements, efficiency at work, and confidence socially are greatly disparate; this can result in decreased levels of comprehension, decreased reading rates, and a greater chance of being misdiagnosed with learning disabilities or attention deficits. On the other hand, students with normal binocular function exhibit enhanced reading fluency, increased academic engagement, and improved overall academic achievement⁴⁰⁻⁴³.

The psychological implications of disorders of binocular vision must also be taken into account³⁹. Those with visible misalignments, as with strabismus, tend to experience social ostracism, low self-esteem, and anxiety. Comparative research shows that those with untreated binocular vision anomalies tend to be more socially withdrawn, participate less in group activities, and have elevated levels of stress relative to those with normal eyes. In addition, research has demonstrated that persons with weak stereopsis avoid sports and activities involving hand-eye coordination,

while those with excellent binocular function perform such activities with greater confidence^{31,35}.

In recent years, advances in vision therapy and assistive technologies have offered promising solutions for enhancing binocular function. Virtual reality (VR)-based rehabilitation has also emerged as a useful tool in vision therapy, offering interactive training exercises to enhance stereopsis and binocular coordination. Comparative studies between traditional vision therapy and VR-based rehabilitation indicate that the latter leads to faster improvement in stereoacuity and binocular function. In addition, the use of prism lenses, digital eye-trackers, and adaptive learning software in schools has been known to help individuals with binocular vision anomalies to better address vision issues^{40,42}.

Despite these developments, heightened awareness among educators, employers, and medical professionals is crucial for early intervention and diagnosis. Schools with routine vision screening programs can identify students at risk of binocular vision disorders early, thereby avoiding academic difficulties and potential misdiagnosis. Employers, particularly those that demand high visual-motor coordination, should provide accommodations and assistive technologies for workers with vision impairments. A comparative study of organisations that have adopted vision-friendly policies and those that have not reveals enhanced productivity, fewer workplace mistakes, and better employee health in the former.

Conclusion

Binocular vision anomalies and reduced stereopsis adversely affect functional performance across educational, occupational, and daily activities. Although modern diagnostic tools and vision-based interventions enable effective management, early detection remains critical. Routine vision screening, increased public and professional awareness and supportive visual environments are essential public-health strategies to reduce disability and optimise functional outcomes in affected individuals.

Acknowledgements

Not Applicable

Funding Statement

There was no Funding.

Conflict of Interest Statement

The authors declare no competing interests.

Data Availability Statement

Not Applicable as it is a review and no data is generated.

References

- 1 Read JCA. Binocular Vision and Stereopsis Across the Animal Kingdom. *Annu Rev Vis Sci*, 7 (2021) 389.
- 2 Guo DD, Wu JF, Hu YY, Sun W, Lv TL, Jiang W, Wu H, Wang XR, Jonas JB & Bi HS. Stereoacuity and related factors: The Shandong children eye study. *PLoS One*. 11 (2016) e0157829.
- 3 Han S, Zhang X, Zhao X, Wang Y, Hao Q, Li R, Huang D, Zhu H & Sun Q. Stereoacuity and Related Factors in Healthy Preschool Children: The Nanjing Eye Study. *Ophthalmic Epidemiol*. 26 (2019) 336.
- 4 Raveendran RN, Bobier WR & Thompson B. Binocular vision and fixational eye movements. *J Vis*. 19 (2019) 9.
- 5 Murphy KM, Roumeliotis G, Williams K, Beston BR & Jones DG. Binocular visual training to promote recovery from monocular deprivation. *J Vis*, 15 (2015) 2.
- 6 Gadia D, Garipoli G, Bonanomi C, Albani L & Rizzi A. Assessing stereo blindness and stereo acuity on digital displays. *Displays*. 35 (2014) 206.
- 7 Dorman R & van Ee R. 50 Years of Stereoblindness: Reconciliation of a Continuum of Disparity Detectors With Blindness for Disparity in Near or Far Depth. *Iperception*. 8 (2017) 2041669517738542.
- 8 Wiebrands M & Woods A & Riddell H. Incidence of stereo blindness in a recent VR distance perception user study. *Electronic Imaging*. 35 (2023) 225.
- 9 Alhusuny A, Cook M, Khalil A, Treleaven J, Hill A & Johnston V. Impact of accommodation, convergence and stereoacuity on perceived symptoms and surgical performance among surgeons. *SurgEndosc*. 35 (2021) 6660.
- 10 Bhattacharya S, Saleem SM & Singh A. Digital eye strain in the era of COVID-19 pandemic: An emerging public health threat. *Indian J Ophthalmol*. 68 (2020) 1709.
- 11 Jackman AH. 3-D cinema: immersive media technology. *GeoJournal*. 80 (2015) 853.
- 12 Zheng F, Hou F, Chen R, Mei J, Huang P, Chen B & Wang Y. Investigation of the Relationship Between Subjective Symptoms of Visual Fatigue and Visual Functions. *Front Neurosci*. 15 (2021) 686740.
- 13 Althomali MM & Leat SJ. Binocular vision disorders and visual attention: Associations with balance and mobility in older adults. *J Aging Phys Act*. 26 (2018) 235.
- 14 Bridge H. Effects of cortical damage on binocular depth perception. *Philos Trans R Soc Lond B Biol Sci*. 371 (2016) 20150254.
- 15 Han SB, Yang HK, Kim J, Hong K, Lee B & Hwang JM. Quantification of Stereopsis in Patients with Impaired Binocularity. *Optom Vis Sci*. 93 (2016) 588.
- 16 Ng CJ, Blake R, Banks MS, Tadin D & Yoon G. Optics and neural adaptation jointly limit human stereovision. *Proc Natl Acad Sci U S A*. 118 (2021) e2100126118.

- 17 Ortiz-Peregrina S, Ortiz C, Casares-López M, Jiménez JR & Anera RG. Effects of cannabis on visual function and self-perceived visual quality. *Sci Rep.* 11 (2021) 1655.
- 18 Metsing TI. The effect of induced blur monocularly and binocularly on stereoacuity. *Afr. Vis. Eye Health.* 82 (2023) a846.
- 19 Zwierko T, Puchalska-Niedbał L, Krzepota J, Markiewicz M, Woźniak J & Lubiński W. The effects of sports vision training on binocular vision function in Female University Athletes. *J Hum Kinet.* 49 (2015) 287.
- 20 Cacho-Martínez P, Cantó-Cerdán M, Carbonell-Bonete S & García-Muñoz Á. Characterization of visual symptomatology associated with refractive, accommodative, and binocular anomalies. *J Ophthalmol.* 2015 (2015) 895803.
- 21 Musch DC, Niziol LM, Gillespie BW, Lichter PR & Janz NK. Binocular Measures of Visual Acuity and Visual Field versus Binocular Approximations. *Ophthalmology.* 124 (2017) 1031.
- 22 Martino F, Castro-Torres JJ, Casares-López M, Ortiz-Peregrina S, Ortiz C & Anera RG. Deterioration of binocular vision after alcohol intake influences driving performance. *Sci Rep.* 11 (2021) 8904.
- 23 García-Muñoz Á, Carbonell-Bonete S, Cantó-Cerdán M & Cacho-Martínez P. Accommodative and binocular dysfunctions: prevalence in a randomised sample of university students. *ClinExpOptom.* 99 (2016) 313.
- 24 Tiwari RP, Barot RK, Sawant V, Wagh U, Maknikar B & Kanetkar M. Clinical Profile of Non Strabismic Binocular Vision Anomalies in MBBS and Nursing Students in a Teaching Hospital: A Cross-sectional Study. *J CliniDiagno Rese.* 16 (2022) 19.
- 25 Devi P, Kumar P, Marella BL & Bharadwaj SR. Impact of Degraded Optics on Monocular and Binocular Vision: Lessons from Recent Advances in Highly-Aberrated Eyes. *SeminOphthalmol.* 37 (2022) 869.
- 26 Iehisa I, Ayaki M, Tsubota K & Negishi K. Factors affecting depth perception and comparison of depth perception measured by the three-rods test in monocular and binocular vision. *Heliyon.* 6 (2020) e04904.
- 27 Verghese P, Tyson TL, Ghahghaei S & Fletcher DC. Depth perception and grasp in central field loss. *Invest Ophthalmol Vis Sci.* 57 (2016) 1476.
- 28 Sverdlitchenko I, Mandelcorn MS, Issashar Leibovtzh G, Mandelcorn ED, Markowitz SN & Tarita-Nistor L. Binocular visual function and fixational control in patients with macular disease: A review. *Ophthalmic Physiol Opt.* 42 (2022) 258.
- 29 Verghese P, Tyson TL, Ghahghaei S & Fletcher DC. Depth perception and grasp in central field loss. *Invest Ophthalmol Vis Sci.* 57 (2016) 1476.
- 30 Mestre C, Otero C, Díaz-Doutón F, Gautier J & Pujol J. An automated and objective cover test to measure heterophoria. *PLoS One.* 13 (2018) e0206674.
- 31 Yaramothu C, Greenspan LD, Scheiman M & Alvarez TL. Vergence endurance test: A pilot study for a concussion biomarker. *J Neurotrauma.* 36 (2019) 2200.
- 32 Bucci MP, Kapoula Z, Bui-Quoc E, Bouet A & Wiener-Vacher S. Saccades and vergence performance in a population of children with vertigo and clinically assessed abnormal vergence capabilities. *PLoS One.* 6 (2011) e23125.
- 33 Vargas V, Radner W, Allan BD, Reinstein DZ, Burkhard Dick H & Alió JL. Methods for the study of near, intermediate vision, and accommodation: an overview of subjective and objective approaches. *SurvOphthalmol.* 64 (2019) 90.
- 34 Ghosh D. Does eye exercises along with physical training helps in achieving better sports performance? Effect of vision therapy on basketball players. *Int. J. Ophthalmol and Optometry.* 6 (2024) 14.
- 35 Aletaha M, Daneshvar F, Mosallaei M, Bagheri A & Khalili MR. Comparison of three vision therapy approaches for convergence insufficiency. *J Ophthalmic Vis Res.* 13 (2018) 307.
- 36 Huang S, Zhong X, Quan C & Zhang M. Non-surgical treatment of strabismus in children: a review of recent advances. *Front Med (Lausanne).* 12 (2025) 1582284.
- 37 Woo Hoi Chi, White Peter & Lai. Ergonomics Standards and Guidelines for Computer Workstation Design and the Impact on Users' Health – A Review. *Ergonomics,* 59 (2016) 464.
- 38 Alrasheed SH & Elmadina AEM. Effect of Binocular Vision Problems on Childhood Academic Performance and Teachers' Perspective. *P J Ophthalmol.* 36 (2020) 163.
- 39 Narayanasamy S, Vincent SJ, Sampson GP & Wood JM. Visual demands in modern Australian primary school classrooms. *ClinExpOptom.* 99 (2016) 233.
- 40 Koulieris George, Bui Bee, Banks Martin & Drettakis George. Accommodation and Comfort in Head-Mounted Displays. *ACM Transac on Graphi,* 36 (2017) 1.
- 41 Aletaha M, Daneshvar F, Mosallaei M, Bagheri A & Khalili MR. Comparison of three vision therapy approaches for convergence insufficiency. *J Ophthalmic Vis Res.* 13 (2018) 307.
- 42 Muhsin ZJ, Qahwaji R, Ghanchi F & Al-Tae M. Review of substitutive assistive tools and technologies for people with visual impairments: recent advancements and prospects. *J Multim Us Interfac.* 18 (2024) 135.
- 43 Li Y, Kim K, Kim K, Norouzi N, Jules J, Bruder G & Welch GF. A Scoping Review of Assistance and Therapy with Head-Mounted Displays for People Who Are Visually Impaired. *ACM Trans Access Comput.* 15(2022) 1.