

## Cytotoxicity and antioxidant activities of methanolic extract of *Marchantia polymorpha* and *Dicranum scoparium* using network pharmacology, molecular docking, and experimental approaches

Shivom Singh<sup>1</sup>, Gaurav Bhadauriya<sup>1</sup>, Anurag Agrawal<sup>2\*</sup>, Vipin Bhati<sup>2</sup>, Shivani Gore<sup>3</sup> and Kajal S. Rathor<sup>4\*</sup>

<sup>1</sup>Department of Environmental Science, <sup>3</sup>Department of Biotechnology, ITM University, Gwalior 474001, Madhya Pradesh, India

<sup>2</sup>Department of Pharmacology, Ram-Eesh Institute of Vocational and Technical Education, Greater Noida 201310, Uttar Pradesh, India

<sup>4</sup>Department of Biotechnology, SMS Govt. Model Science College, Gwalior 474009, Madhya Pradesh, India

Received 31 March 2024; revised received 23 October 2024; accepted 29 October 2024

### Supplementary Figures




Fig. 1 — Common target identifies for gallic acid.




Fig. 2 — Top 10 genes network of gallic acid.




Fig. 3 — Common targets were identified for benzoic acid.




Fig. 4 — Top 10 gene network benzoic acid.




Fig. 5 — Common targets were identified for lupeol.




Fig. 6 — Top 10 gene network lupeol.




Fig. 7 — Common targets were identified for quercetin.




Fig. 8 — Top 10 gene network with quercetin.