



A review of the nutritional potential of edible snails: A sustainable underutilized food resource

Mahendra Baghele¹, Shubhi Mishra¹, Victor Benno Meyer-Rochow^{2,3}, Chuleui Jung^{2,4} and Sampat Ghosh^{2*}

¹Department of Life Sciences, Sardar Patel University, Balaghat, Madhya Pradesh 481331, India

²Agriculture Science and Technology Research Institute, Andong National University, Andong, Gyeongsangbuk 36729, Republic of Korea

³Department of Genetics and Ecology, Oulu University, Oulu 90140, Finland

⁴Department of Plant Medicinals, Andong National University, Andong, Gyeongsangbuk 36729, Republic of Korea

Received 24 March 2021; revised received 04 November 2022; accepted 24 November 2022

Despite the use of snails as food in many different communities in the world, the consumption of snail meat is still far from being appreciated. The authors compiled and reviewed the nutrient composition of edible snail meat based on the available scientific literature. The protein, fat, and ash content of snail meat were found to be 14.0, 1.4 and 2.1% on the basis of fresh matter respectively. Comparable protein content with conventional foods of animal origin along with the presence of all essential amino acids and less carbon footprint makes the snail meat a sustainable food source. Moreover, less fat content but higher proportion of polyunsaturated fatty acids, and considerable presence of minerals with nutritional importance especially calcium, zinc, and iron could exhibit human nutritional benefits. However, the establishment of heliculture facilities is a necessity in order to support this sustainable food resource and also enhance the economic condition of certain local areas.

Keywords: Amino acids, Angiostrongyliasis, Food safety, Food security, Minerals, Protein

IPC code; Int. cl. (2021.01)-A23J, A61K 36/00

Introduction

According to the recent report of the global hunger index¹, 828 million people are suffering from undernourishment (defined as a lack of sufficient calories in the food) and inadequate nutrition (defined as a lack of nutrients in the diet). Sub-Saharan Africa and South Asia are regions with the highest hunger and undernutrition levels in the world¹. At least 2 billion people are suffering from micronutrient deficiency which is often termed “the hidden hunger”², but at the same time about 2 billion people suffer from overweight or obesity. In the context of a still expanding global population and the unanticipated climate change, we as humans are forced to explore alternative sustainable food sources to feed the people in future.

We need to increase the production of nutrient-rich food without harming the environment or increasing greenhouse gas emissions, or demanding to provide additional water, but with an aim to achieve low feed conversion ratios and to intensify farming systems

without laying off farm workers. To understand the linkages between sustainability, productivity or yield of food not only the unit area of land has to be considered but inputs like irrigation needs, fertilizers, insecticides, herbicides, machinery, fuel costs, emission of greenhouse gases (GHG) etc. must not be overlooked either. It was first suggested by Meyer-Rochow to use unconventional (alternative) foods, like for instance edible insects to ease global food shortages and to involve the FAO as well as WHO³. The idea slowly gained acceptance and has led to the current interest in entomophagy⁴. Algae as a food source has also received some attention in the last few decades⁵, but snails as a candidate to augment the spectrum of available food items has been given a comparatively poor attention in the agriculture sector. This is somewhat surprising because a variety of snail species are even now commonly consumed by members of numerous communities around the world. In fact, the use of snails as food has a long history and in the Mediterranean region has been known from prehistoric times⁶ to this day⁷.

According to the estimates of FAOSTAT⁸, there is an increasing trend to produce food based on snails (marine

*Correspondent author
E-mail: sampatghosh.bee@gmail.com

species not included) year by year (Fig. 1). On average, 15% of the snails come from snail-breeding units, while the remaining 85% are harvested from Nature⁹. Snail farming, known as ‘Heliciculture’, is not at all a new entrant in the agriculture sector, but initiatives to explore this food resource have been limited and the resource clearly remains as under-utilized.

In the era of globalization, we tend to lose traditional foods with the result of nutrient deficiencies and negative health consequences¹⁰. Most often the interruption of a regular supply plays a critical role in the declination of traditional foods. Halwart, reported that there was no tradition of snail farming and marketing in south-east Asia although rural people of different parts of this region preferred to eat native snails¹¹. In Japan, snail farming became popular in the 90s, and many governmental and non-governmental livelihood projects were established to support a backyard cottage industry¹¹. Snails have also been used as feed for broiler chickens, ducks, and pigs¹¹⁻¹³. However, because of an assumed high pest potential, farming systems did not continue to use snails much as animal feed and, to cite an example, Vietnam banned snail farming in mid-1992. In many parts of India, local people do collect edible terrestrial as well aquatic snails from the forest and water bodies and consume them¹⁴ (personal observations of SG). Recently some small scale snail farms have been developed (personal communication of SG).

In a survey on bushmeat in Nigeria, Martin placed snails in a separate category from bushmeat as they were very common (especially the giant African snail *Archachatina marginata*) particularly in the forested areas where people did not consider them as bushmeat¹⁵. For the last few decades Africa, and in particular western Africa, is experiencing a rapid growth of the number of snail farms with the adequate help of governmental as well as non-governmental agencies¹⁶⁻¹⁸. Snail farming has also received attention in developed country like Korea. Ghosh *et al.*, described the farming system of *Pomacea canaliculata*, as edible snail, as it is practiced as mini-livestock in Korea¹⁹. From 1995 to 2010, Italian snail production and consumption increased almost four times and the annual sale in 2013 was worth about 265 million Euros²⁰. Allen estimated that the consumption of edible snail *Theba pisana* accounts for approximately 4000 tons per year^{7,21}. Snail farming is characterized by not involving labour-intensive work, or requiring high capital investment, or sophisticated technical and professional knowledge, and this makes it suitable for developing and/or underdeveloped areas. However, in order to appreciate the nutritional advantages of the rearing and consumption of snails, it is essential to examine and highlight the nutrient composition of the snails. The primary aim of this review has therefore been to evaluate the nutrient content of the edible species.




Fig. 1 — Global production of snails (marine species not included) with the year (Source: FAOSTAT, 2021).

Edible snail species in the world

Based on the available scientific literature, the authors compiled an inventory of edible snails around the globe and presented it in Table 1. Although there

are undoubtedly many more, we focused on those species that are widely popular and have, to some extent, been scientifically studied with regard to chemical composition, biology, and breeding. Such

Table 1 — Inventory of edible snail and bivalve species around the world

Scientific name	Geographical location	Purpose	Reference
Africa			
<i>Helix aspersa</i>	Morocco	Food	50
<i>Limicolaria aurora</i>	Uyo main market, Akwa Ibom state, Nigeria	Food	51
<i>Archachatina marginata</i>	Ikere-Ekiti market, Ondo state, Nigeria	Food	52
<i>Archatina</i>			
<i>Limicolaria</i>			
<i>Achatina achatina</i>	Ghana	Food	53
<i>Limicolaria</i> sp.	Oba market (Ado-Ekiti), Ekiti state, Nigeria	Food	54
<i>Archatina archatina</i>			
<i>Archachatina amarginata</i>			
<i>Lymnaea stagnalis</i>	Kuto market in Abeokuta, Ogun state, Nigeria	Food	55
<i>Archachatina marginata ovum</i>	King's market, Ado Ekiti, Ekiti state, Nigeria	Food	56
<i>Archachatina marginatasaturalis</i>			
<i>Achatina achatina</i>			
<i>Limicolaria</i> sp.			
<i>Archachatina marginata ovum</i>	Wild collection from Calabar, Odukpani and Ugep;	Food	57
<i>Archachatina marginatasaturalis</i>	Nigeria		
<i>Limicolaria</i> sp.			
<i>Lanistes varicus</i>			
<i>Nucella lapillus</i>			
<i>Archachatina marginata</i>	Markets in Ibadan metropolis, Oyo state, Nigeria	Food	58
<i>Achatina achatina</i>			
<i>Achatina fulica</i>			
<i>Achatina fulica</i>	Uyo, Akwa Ibom state, Nigeria	Food	59
<i>Limicolaria</i> sp.			
<i>Helix pomatia</i>			
<i>Helix trunculus</i>	Sfax city fish market, Tunisia	Food	60
<i>Archachatina marginata</i>	Minna, Niger state, Nigeria	Food	61
<i>Archachatina marginata</i>	Farm at Rivers State University of Education, Ndele campus, Nigeria	Food	62
<i>Achatina achatina</i>			
<i>Limicolaria flammea</i>	Farm in Kumasi, Ghana	Food	63
<i>Archachatina marginata</i>			
Agboville, Côte d'Ivoire			
	Itoku market, Abeokuta, Ogun state, Nigeria	Food	64
	Oje market, Oyo state, Nigeria	Food	65
	Oyingbo market, Lagos state, Nigeria		
	Oja Oba market, Osogbo, Osun state, Nigeria		
	Oja Oba market, Akure, Ondo state, Nigeria		
<i>Archachatina marginata</i>	Oyo state, Nigeria	Food	66
<i>Achatina fulica</i>	Littoral (Wouri), Center (Lekie) and West (Santchou),	Food	67
<i>Archachatina marginata</i>	Cameroon		
Asia			
<i>Pila scutata</i>	Malaysia	Food	68
<i>Bellamyiaingallsiana</i>			
<i>Ampullarius insularis</i>	Jepllabuk-do, Korea	Food	69
<i>Achatina fulica</i>			
<i>Pomacea canaliculata</i>			
	Thailand	Food	11
	Vietnam		
	Taiwan		
	Philippines		
<i>Achatina fulica</i>	Junchun Food Co., Kimhae, Korea	Food	70
<i>Pila globosa</i>	29 tribal groups in Bangladesh	Food	71,72
Apple snail	Japan	Food	73
<i>Helix pomatia</i>	Local processing company in Cukurova, Turkey	Food	74

(Contd.)

Table 1 — Inventory of edible snail and bivalve species around the world (*Contd.*)

Scientific name	Geographical location	Purpose	Reference
<i>Semisulcospira gottschei</i>	Korea	Food	75
<i>Achatina fulica</i>	China	Food	76
<i>Pomacea canaliculata</i>			
<i>Bellamyia bengalensis</i>	North East India	Food	77
<i>Bellamyia bengalensis f. typica</i>			
<i>Bellamyia bengalensis f. balteata</i>			
<i>Bellamyia dissimilis</i>			
<i>Pila globosa</i>			
<i>Pila scutata</i>			
<i>Pila theobaldi</i>			
<i>Pila virens</i>			
<i>Brotia costula</i>			
<i>Bellamyia sp.</i>	Northern part of Bihar (Khagaria, Saharsa, Medhepura,	Ethnomedicinal	23
<i>Bellamyia bengalensis</i>	Purnea), India	uses	
<i>Pila scutata</i>			
<i>Lamellidens sp.</i>			
<i>Parreysia sp.</i>			
<i>Lamellidens sp.</i>			
<i>Semisulcospira coreana</i>	Korea	Food	78
<i>Semisulcospira gottschei</i>			
<i>Semisulcospira forticosta</i>			
<i>Semisulcospira libertina</i>			
<i>Semisulcospira tegulata</i>			
<i>Helix aspersa</i>	Asian Istanbul, Turkey	Food	79
Viviparid snail	Vietnam	Food	80,27
<i>Chamberlainia hainesiana</i>	Thailand, Vietnam	Food	
<i>Brotiacostula</i>	Wild (pond, paddy field, wetland) and market,	Food	81
<i>Bellamyia bengalensis</i>	Lembucherra, Tripura, India		
<i>Bellamyia dissimilis</i>			
<i>Pila globosa</i>			
<i>Helix aspersa</i>	Tribal people of Chhindwara (Gond, Bharia, Mawasi,	Ethnomedicinal	24
	Ahir), Madhya Pradesh, India	uses	
<i>Pila sp.</i>	Tribal belt of Gondwanavindhyan region, Madhya	Ethnomedicinal	25
	Pradesh, India	uses	
<i>Pomacea canaliculata</i>	Farm in Andong, Korea	Food	82
<i>Pila globosa</i>	Consumed by Muda tribe, Jharkhand, India	Food	83
Unidentified snails species	Local markets (collected from wild areas like water	Food	Personal experience
	bodies) in northern part of West Bengal, India		of SG
Unidentified snails species	Consumed by Baiga tribe, Madhya Pradesh, India	Food	14
	Australia		
<i>Cookia sulcata</i>	Te Oka Bay, Canterbury, New Zealand (consumed by	Food	84
	Maori people)		
<i>Turbo militaris</i>	Woolgoolga, Coffs harbour, Australia	Food	85
<i>Lunella undulata</i>			
<i>Lunella torquata</i>			
	Europe		
<i>Helix adpersa</i>	Europe	Food and	22,86
<i>Helix pomatia</i>		Therapeutic	
<i>Helix lucorum</i>		uses	
<i>Helix aperta</i>			
<i>Helix cincta</i>			
<i>Otala sp.</i>			
<i>Eobania vermiculata</i>			
<i>Helix pomatia</i>	Farm and wild, Latvia	Food	87
<i>Lymnaea sp.</i>	Europe	Therapeutic uses	22
<i>Thebapisana</i>	Portugal	Food	7
<i>Otalalactaea</i>	Portugal	Food	
	South America		
<i>Helix aspersa maxima</i>	Brazil	Food	88,89

species include the snails *Achatina* sp., *Archachatina marginata*, *Limicolaria* sp., *Lanistes varicus*, *Nucella lapillus*, *Lymnaea stagnalis*, *Helix* sp., *Pila* sp., *Semisulcospira* sp., *Pomacea canaliculata*, *Bellamya* sp., *Theba pisana* and *Otala lactea*. *Escargots* (mainly *Helix pomatia* but some other species as well) are considered a delicacy in France and some other European countries. A systematic review of the literature reveals that the consumption of snails as a planned part of the diet is more common in Africa and Asia, although the acceptance varies to a great extent from region to region.

Although these species are used primarily as food, ethnomedicinal uses as reviewed by Meyer-Rochow are also common and involve a wide variety of slugs and snails in connection with dermatoses, gastro-enteric troubles, coughs, colds, heart ailments and even bone fractures²². Consumption of snail soup or curry containing, for example, *Bellamya* sp., *Pila* sp., *Lamellidens* sp., or *Parreysia* sp. is common among the people of the Kosi River basin of Bihar (India) for treating asthma, arthritis, rickets, cardiac disorders, giddiness, night blindness, etc.²³. An ethnozoological use of the garden snail (*Helix aspersa*) for the treatment of tuberculosis by the indigenous people of the Chindwara District of Madhya Pradesh (India)²⁴, the use of *Pila* sp. snails for treating weakness and paralysis by the tribal people and communities of the Gondwana Vindhyan region of Madhya Pradesh (India)²⁵ and the use of snails for bone strength, for treatment of body ache, joint pain, etc. by the Baiga people of Baihar tehsil of Balaghat district of Madhya Pradesh (India)¹⁴ are examples of therapeutic uses of snails in India.

Edible snails belong to the molluscan class Gastropoda and families like Achatinidae, Helicidae, Ampullariidae, Viviparidae, Thiaridae, Planorbidae, Pachychitidae, Turbinidae are represented. In general, a generic freshwater snail is dioecious, while those belonging to the subclass Pulmonata are hermaphrodites²⁶. Even in the case of the hermaphroditic species mating between two individuals occur followed by the deposition of eggs often covered with a protective gelatinous coat. Although developmental duration and life cycles are species-specific, under favourable conditions eggs usually hatch after about 2 weeks to one month. Thiarid (family Thiaridae) females are parthenogenic²⁶ and Ampullarids (family Ampullariidae), commonly known as 'apple snails',

are mostly amphibious with a mantle cavity containing organs that function as gills and lungs. *Pila* is a common and widely distributed genus in tropical Asia, but *Pomacea canaliculata* is an example that was introduced to many Asian countries from the USA and South America and became established^{11,27}. The Achatinidae is a family containing terrestrial pulmonate gastropods from Africa, such as the genera *Achatina*, *Limicolaria*, and *Archachatina*. Viviparidae are often known as river snails and contain live-bearing species in which the females do not lay eggs, but retain them in their body.

Nutrient composition of edible snails

Moisture content

The moisture content of edible snail species varies within a range of 65.8 to 89.2% (Fig. 2 and Table 2). Fresh snails are generally used in preparations that include boiling, smoking, and using them for curries. In France, *Escargots-in-garlic-and-parsley-butter* is a famous snail dish; other worthwhile snail cuisines are the *Gaeng-Kui-Hoi-Khom* fresh water snail in red curry in Thailand, the *Tharoi Thongba* river snail curry in Manipur (India), the *Óchuong* stir-fried snail and *Bún Óc* in Vietnam, the *Lumache alla Romana* in Rome (Italy), *Cargols* in Spain, *Caracóis* stewed snail in Portugal, and *Kohlibourbouristi* pan-fried snail in Greece. In general, the high moisture content of snails makes their meat more susceptible to microbial contamination²⁸. Therefore in order to increase the




Fig. 2 — Proximate composition (% of fresh weight basis) of edible snail species. Number indicates the mean value of corresponding proximate composition i.e., moisture, protein, fat, ash [Graphical representation was plotted based on Table 2].

Table 2 — Proximate composition of edible snail species

Scientific name	Proximate composition (% Fresh weight basis)						Reference
	Moisture	Protein	Lipid	Fibre	Ash	NFE/Carbohydrate	
	Land						
<i>Achatina</i> sp.	89.20	9.90	1.40	--	2.10	4.40	90
<i>Helix aspersa</i> (small, 2.5 g)	83.30	12.00	0.70	--	2.70	0.50	91
<i>Helix aspersa</i> (large, 3.5 g)	87.60	9.90	0.50	--	1.20	0.40	
<i>Achatina fulica</i> (Gray)	82.18	11.53	0.91	--	1.39	--	70
<i>Achatina fulica</i> (Yellow)	82.36	13.42	1.28	--	1.29	--	
<i>Achatina fulica</i> (White)	81.20	13.69	0.97	--	1.25	--	
<i>Limicolaria aurora</i> *	71.24	14.78	2.79	--	3.38	7.79	51
<i>Archachatina marginata</i>	76.63	20.76	1.20	ND	1.42	--	52
<i>Archatina</i> sp.	79.89	14.52	4.24	ND	1.34	--	
<i>Limicolaria</i> sp.	77.90	17.51	4.30	ND	1.12	--	
<i>Helix aspersa aspersa</i>	83.70	10.50	1.60	--	1.50	--	92
<i>Helix aspersa maxima</i>	85.80	8.10	1.50	--	1.20	1.70	
<i>Helix lucorum</i>	84.00	10.80	1.50	--	1.40	--	
<i>Helix pomatia</i>	84.20	10.70	1.10	--	1.80	--	
<i>Helix aspersa maxima</i> ***	79.85	9.50	2.66	--	0.92	--	89
	79.05	10.47	1.51	--	0.65	--	
	77.47	11.31	0.45	--	0.76	--	
	78.41	12.56	0.63	--	0.65	--	
<i>Helix aspersa</i>	80.80	16.35	0.41	--	1.89	--	74
<i>Archachatina marginata</i> (ovum) Pfeiffer	76.56	20.56	1.38	ND	1.44	0.01	56
<i>Archachatina marginata</i> (saturalis) Philippi	76.67	20.34	1.23	ND	1.40	0.37	
<i>Achatina achatina</i>	77.54	19.27	1.43	ND	1.34	0.42	
<i>Limicolaria</i> sp.	78.68	18.66	1.17	ND	1.35	0.15	
<i>Archachatina marginata ovum</i> *	80.78	16.23	0.86	0.10	0.77	1.26	57
<i>Archachatina marginata saturalis</i> *	80.30	15.95	0.79	0.20	0.89	1.89	
<i>Limicolaria</i> sp.*	78.64	15.33	0.80	0.21	1.50	3.31	
<i>Archachatina marginata</i>	73.67	19.53	2.44	--	2.56	1.80	58
<i>Achatina achatina</i>	75.28	17.20	2.21	--	2.33	2.98	
<i>Achatina fulica</i>	79.28	10.08	1.61	--	1.78	7.25	
<i>Limicolaria</i> sp.	84.91	5.86	1.05	--	1.27	6.91	
<i>Helix pomatia</i>	79.57	15.44	1.18	0.00	1.26	--	59
<i>Helix</i> sp.	83.20	8.64	0.57	0.004	1.04	6.55	93
<i>Helix aspersa</i>	82.50	12.87	0.58	--	1.07	4.99	79
<i>Archachatina marginata</i> **	73.95	20.22	2.09	1.01	1.73	1.09	65
	79.67	15.77	1.46	0.79	1.35	0.86	
	78.85	16.42	1.60	0.82	1.40	0.92	
	77.41	12.54	1.73	0.88	1.51	0.95	
	77.00	17.88	1.77	0.89	1.53	0.94	
<i>Archachatina marginata</i> (Adult)	79.25	17.22	1.46	--	1.34	--	94
<i>Archachatina marginata</i> (Grower)	79.93	16.30	1.35	--	1.24	--	
<i>Archachatina marginata</i> (Snaillet)	80.39	15.83	1.24	--	1.18	--	
<i>Otala lactea</i> (boiled)	83.50	9.60	1.26	--	1.38	0.28	7
<i>Otala lactea</i> (cooked)	78.60	12.90	1.54	--	3.45	0.32	
<i>Theba pisana</i> (boiled)	83.20	11.40	2.33	--	1.11	0.21	
<i>Theba pisana</i> (cooked)	77.50	13.40	3.23	--	3.05	0.35	
	Freshwater						
<i>Pila globosa</i>	85.50	8.27	0.73	0.03	2.60	2.90	93
<i>Bellamyia bengalensis</i>	82.10	8.97	0.98	0.04	3.64	4.31	
<i>Melaniatu berkulata</i>	74.60	12.36	1.79	0.05	3.68	7.57	
<i>Anisus convexiusculus</i>	75.70	12.93	0.97	0.04	4.61	5.79	
<i>Brotia costula</i>	69.86	12.91	0.82	ND	7.28	9.12	81
<i>Bellamyia bengalensis</i>	65.80	13.14	0.96	ND	8.11	11.97	
<i>Bellamyia dissimilis</i>	65.80	11.18	0.96	ND	8.15	11.46	
<i>Pila globosa</i>	73.80	15.59	1.15	ND	3.82	5.62	

(Contd.)

Table 2 — Proximate composition of edible snail species (*Contd.*)

Scientific name	Proximate composition (% Fresh weight basis)						Reference
	Moisture	Protein	Lipid	Fibre	Ash	NFE/Carbohydrate	
	Marine						
<i>Turbo militaris</i>	73.08	16.19	5.57	--	2.14	3.02	85
<i>Lunella undulata</i>	70.83	18.49	5.20	--	1.97	3.51	
<i>Lunella torquata</i>	68.50	18.03	8.46	--	2.10	2.92	

*Calculated from obtained data on the dry weight basis to fresh weight basis
 **Five values were obtained for the samples collected from five regions
 ***Four values were obtained for each component because of four different feed treatments
 ND= Not determined




Fig. 3 — Comparative account of a) protein content (g/100 g fresh weight basis) of edible snails and conventional foods of animal origin; and b) fat content (g/100 g fresh weight basis) of edible snails and conventional foods of animal origin. [Data other than edible snails were obtained from USDA database <https://fdc.nal.usda.gov> accessed 10th February 2021].

shelf life of snails, drying prior to processing them further can be useful.

Protein and amino acids

The protein content of snail meat was found to be within the range of 5.9 to 20.8% with an average value of 14.0% on the basis of fresh matter (Fig. 2 and Table 2). The protein content of edible snails was therefore comparable to that of conventional animal meats (Fig. 3a). However, if converted on the dry weight basis using an average moisture content of 78.7%, the protein content reaches approximately 65.7%, which is higher than the protein contents of conventional protein sources. The nutritive quality of a protein depends on its amino acid composition. Fig. 4 represents the amino acid compositions of edible snails (Table 3). For several species, tryptophan was not determined presumably due to the acid hydrolysis process. Cysteine and methionine amounts are also generally not unambiguously identifiable due to the acid hydrolysis process in the analysis. Apart from the methodological concerns, the average proportion of essential amino acids was found to be about 36.5% while the nonessential amino acids accounted for 51.5%. Of course, the proportion will




Fig. 4 — Compositional distribution of amino acids on the basis of % of protein of edible snails. Number indicates the mean value of corresponding individual amino acid [Graphical representation was plotted based on Table 3].

be enhanced if cysteine and tyrosine are included as conditionally essential amino acids and arginine as essential for children. Among the essential amino acids, leucine (5.4-11.5%) and lysine (5.0-8.4%) are found predominant while glutamic acid (1-17.3%)

Table 3 — Amino acid composition of edible snails

Scientific name	Amino acid composition (g/100 g protein)																Reference		
	Val	Leu	Lys	Ile	Thr	Met	Cys	Trp	His	Phe	Tyr	Ser	Arg	Pro	Asp	Glu		Gly	Ala
<i>Achatina fulica</i> (Gray)	6.40	7.19	5.33	5.35	4.91	2.46	1.43	0.68	2.09	4.61	2.82	4.37	7.57	7.11	8.37	13.59	6.38	3.88	70
<i>Achatina fulica</i> (Yellow)	6.46	6.64	4.95	4.83	4.93	3.63	1.44	0.53	3.08	4.45	2.31	5.13	7.00	5.57	8.73	14.37	6.27	4.39	
<i>Achatina fulica</i> (White)	6.84	6.64	5.00	4.85	5.00	3.45	1.44	0.65	2.95	4.46	2.28	4.94	7.05	5.69	8.09	14.61	6.23	4.38	
<i>Limicolaria aurora</i>	7.60	11.50	7.20	7.30	3.90	1.40	0.90	ND	3.70	6.70	5.90	4.00	0.30	5.60	7.80	13.50	6.50	6.90	51
<i>Limicolaria</i> sp.	4.12	6.90	8.28	5.24	1.80	1.81	0.48	ND	3.56	4.37	2.58	4.05	6.74	3.99	7.61	1.01	5.06	3.98	54
<i>Archatina archatina</i>	3.28	5.58	6.06	4.46	2.24	1.37	0.87	ND	2.96	3.84	2.48	3.17	6.31	3.01	6.94	11.10	5.16	2.42	
<i>Archachatina marginata</i>	7.11	8.11	5.74	3.87	2.76	2.00	1.09	ND	4.70	4.73	2.93	3.33	5.96	3.61	7.27	14.4	4.57	5.19	
<i>Hexaplex trunculus</i>	4.72	6.96	8.35	4.81	1.93	1.99	0.36	ND	3.92	4.19	3.17	4.36	7.48	3.01	7.90	1.23	6.35	3.56	60
<i>Helix aspersa</i> *	6.34	5.42	6.39	4.12	4.00	3.78	3.47	ND	2.25	3.22	5.29	9.22	5.98	2.84	8.84	12.46	6.94	9.44	79
<i>Cookia sulcata</i> (small)	2.50	5.60	6.00	2.60	3.40	0.30	0.20	4.20	1.30	2.40	2.30	3.80	10.10	5.10	7.00	13.80	9.80	4.50	84
<i>Cookia sulcata</i> (big)	2.50	6.50	5.70	2.50	3.50	0.30	0.30	5.30	1.40	2.40	2.30	3.80	10.20	4.80	6.90	14.10	9.60	4.40	
<i>Pomacea canaliculata</i>	4.30	8.20	6.80	4.10	4.70	1.00	0.80	0.20	1.60	4.30	4.50	4.90	9.10	4.70	8.50	17.30	5.80	5.90	82

*Calculated based on obtained values on the basis of total amino acids as protein

ND = Not determined

followed by aspartic acid (6.9-8.8%) was most abundant among the nonessential amino acids.

Currently, on average animal proteins contribute 25% of the protein intake worldwide, providing 18% of the calories²⁹. From the nutritional point of view, animal meat contains high-quality protein and highly bioavailable micronutrients like vitamin A, vitamin B12, riboflavin, iron, zinc, calcium, etc.³⁰. The consumption of meat in developed countries is much higher than its consumption in underdeveloped as well as developing regions of the world²⁹. However, the socio-economic changes like economic development, increased urbanization and globalization are leading to a transition in dietary patterns particularly in low and medium-income countries and increasing demand for animal protein is being observed³¹. At present, ruminants, chickens and pigs are contributing 96% to the global supply of animal protein, while aquaculture is growing fast³². However, not only the amount of protein available to the underprivileged societies is of concern, but to improve the production system and the distribution of food and at the same time guarantee sustainability is the challenge of the day. Forte *et al.*, demonstrated that the snail carbon footprint amounted to 0.7 kg CO₂ eq. per kg of edible meat²⁰, which is far less than the amount generated by the conventional food sources of animal origin like chicken (1.8-5.2), pork (3.6-8.9), beef (11-51)³³, and even the newly advocated mealworm food source with its 2.7 kg CO₂ eq. per kg³⁴. Unlike cattle, the carbon footprint of snails is not affected by biogenic enteric methane emission but mainly derived from the feed crop cultivation and processing stages²⁰.




Fig. 5 — Distribution of different categories of fatty acids, i.e. Saturated fatty acids (SFA), Monounsaturated fatty acids (MUFA) and Polyunsaturated fatty acids (PUFA) on the basis of % of total fatty acids of edible snails. Number indicates the mean value of corresponding category of fatty acid [Graphical representation was plotted based on Table 4].

Fat and fatty acid composition

Fat content of snail meat was found to be within the range of 0.4 to 8.5% with an average value of 1.4% on the basis of fresh matter (Table 2). Fat content of edible snails was found lower than that of conventional foods of animal origin (Fig. 3b). Data on snail meat fatty acid profiles available from the literature have been compiled in Fig. 5 and Table 4. Based on their degree of saturation, fatty acids are categorized into three groups, i.e. saturated fatty acids (SFA), mono-unsaturated fatty acids (MUFA) and poly-unsaturated fatty acids (PUFA). With the exception of a few cases, the PUFA content of edible

Table 4 — Fatty acid composition mentioning selective predominant fatty acids and three categories of fatty acids i.e. SFA, MUFA, PUFA (% of total fatty acids) of edible snails

Scientific name	Fatty acid composition (%)									Reference
	Myristic	Palmitic	Stearic	Total SFA	Oleic	Total MUFA	Linoleic (n-6)	Linolenic (n-3)	Total PUFA	
<i>Helix pomatia</i>	0.42	10.29	16.38	37.87	14.70	19.65	13.56	1.87	25.83	74
<i>Helix aspersa maxima</i>	0.57	10.11	13.47	26.26	20.42	23.82	19.37	1.04	49.92	89
	0.42	7.68	14.51	24.86	18.61	22.68	19.74	1.10	52.46	
	0.52	8.11	12.32	23.06	18.32	21.63	24.53	1.34	55.31	
	0.65	6.72	12.59	22.20	16.73	20.74	22.70	1.32	57.06	
<i>Helix aspersa</i>	ND	7.28	17.37	28.76	13.57	20.66	17.53	5.63	34.38	79
<i>Hexaplex trunculus</i>	1.23	6.21	6.01	33.44	3.08	9.84	0.83	3.07	58.37	60
<i>Melanopsis praemorsa</i> *	5.66	20.04	8.37	39.00	10.36	21.30	16.33	8.87	39.98	95
	2.35	16.43	8.65	31.12	24.20	40.28	9.37	11.00	28.15	
	5.86	26.48	10.70	46.87	12.60	21.92	12.11	10.72	31.68	
	10.58	33.01	7.27	57.61	10.26	17.33	7.02	5.07	25.50	
<i>Cookia sulcate</i> (Small)	2.30	27.30	8.10	46.60	8.80	14.90	2.30	0.50	32.30	84
<i>Cookia sulcate</i> (Large)	2.30	2.40	9.60	42.20	8.00	10.80	2.10	1.00	36.30	
<i>Assyriellaescheriana</i> †	0.58	10.16	10.30	23.05	12.68	14.97	20.45	8.42	62.38	96
<i>Assyriellaguttata</i> †	1.85	9.61	12.33	26.95	15.06	16.14	18.54	6.24	56.80	
<i>Lunellatorquata</i>	0.55	23.01	5.68	39.95	8.21	14.45	1.60	0.84	45.59	85
<i>Lunellaundulata</i>	0.96	21.62	6.59	38.61	8.41	14.83	2.82	2.33	46.58	
<i>Turbo militaris</i>	0.13	22.12	5.45	39.83	7.89	14.05	2.92	1.87	45.13	
<i>Pomacea canaliculata</i>	3.10	20.50	9.00	39.50	9.10	17.70	20.60	0.90	42.70	82
<i>Helix lucorum</i>	0.53	9.66	6.75	21.70	22.02	26.53	21.76	10.08	51.77	97
<i>Eobania vermiculata</i>	0.30	10.49	7.76	22.68	19.07	21.77	37.67	7.31	55.56	
<i>Helix pomatia</i> (Farm A)††	0.47	10.16	14.31	34.84	13.83	19.55	17.95	2.47	39.68	98
	0.52	11.24	15.51	37.98	15.57	19.91	16.95	1.57	34.14	
	0.54	11.32	15.87	39.26	15.69	20.06	16.69	1.42	32.11	
<i>Helix pomatia</i> (Farm B)††	0.47	10.28	14.37	35.21	13.65	19.83	17.87	2.54	38.78	
	0.54	11.39	15.64	38.38	15.63	20.00	16.59	1.40	32.63	
	0.56	11.45	15.84	38.40	15.70	20.16	16.47	1.41	30.89	
<i>Cornuaspersum maxima</i> (Farm A)††	0.34	8.36	12.18	28.72	20.36	24.22	32.42	3.50	44.58	
	0.38	10.13	13.74	32.69	23.47	24.78	30.42	1.82	38.58	
	0.38	10.31	13.66	33.49	23.50	27.88	30.43	1.45	38.00	
<i>Cornuaspersum maxima</i> (Farm B)††	0.35	8.27	12.23	28.32	26.64	30.91	27.42	3.70	39.81	
	0.40	10.26	13.83	32.88	30.79	31.48	24.55	2.80	31.54	
	0.43	10.34	13.91	33.80	30.80	35.59	25.51	2.70	31.57	
<i>Cornuaspersumaspersum</i> (Farm A)††	0.33	8.93	12.28	29.35	19.56	23.89	34.34	3.70	46.65	
	0.37	10.17	13.87	32.87	23.18	24.25	31.84	2.22	40.11	
	0.38	10.5	13.96	33.92	23.40	28.13	31.71	2.21	39.34	
<i>Cornuaspersumaspersum</i> (Farm B)††	0.35	8.66	12.89	29.54	25.81	29.85	27.40	3.85	41.30	
	0.41	10.22	13.77	32.90	28.74	33.55	25.12	2.86	33.99	
	0.44	10.51	13.94	33.93	29.02	33.72	25.02	2.44	33.09	
<i>Thebapisana</i> **	0.73	12.63	5.41	20.72	28.83	31.27	18.78	7.64	48.10	99
	1.06	15.75	6.31	25.32	28.83	30.49	21.35	8.87	44.19	
<i>Cornuaspersum</i> **	0.76	16.02	7.72	26.97	23.79	25.81	22.15	15.78	47.22	
	0.59	13.24	7.24	23.01	29.95	33.53	19.07	15.14	43.56	
<i>Eobania vermiculata</i> **	0.81	14.63	7.66	25.58	26.03	27.86	21.94	12.40	46.56	
	0.63	13.31	7.59	23.35	29.71	32.24	19.20	15.53	44.41	

* fatty acid content of the species was investigated in different seasons like autumn, winter, spring and summer (from the upper row downwards respectively)

† these species are not eaten

†† fatty acid profile was investigated in three forms i.e. raw, cooked and frozen (from upper row downwards respectively)

** the species was examined on the raw (upper row) and boiled (lower row) condition

snails is predominating and followed by SFA and MUFA contents. SFA content varied within the range of 20.7 to 57.6%, while MUFA and PUFA accounted for 9.8-40.3 and 25.5-62.4%, respectively. Statistical analysis (one-way ANOVA) with the available values for different categories of fatty acids reveals a significant difference among the three categories of fatty acids ($df = 2, 132; F = 57.435, p = 0.000$). Fatty acids play many different critical roles in the human body, being structural components of membrane lipids, precursors of molecules like eicosanoids, prostaglandins, and essential components of the metabolism. Saturated fatty acids, in general, are not desirable, because of their linkage with an elevation of cholesterol and thus cardiovascular disease. Lauric, myristic, and palmitic acids contribute comparatively more to a higher level of cholesterol³⁵, but stearic acid has been reported to lower the cholesterol level³⁶⁻³⁸. Among the SFAs, palmitic and stearic acid were the most abundant in the snail's fat, but oleic acid was the dominant component of the MUFAs. However, the distribution of the fatty acids belonging to PUFA category varied so widely that generalizations at this stage are not possible.

Mineral contents

Minerals contents of edible snails are shown in Fig. 6(a-b) and Table 5. Calcium, an essential macro-mineral, is found to be one of the most abundant minerals in edible snails. In humans, the majority of calcium by virtue of its phosphate salt is present in bones and teeth. It plays vital roles in several enzyme-mediated processes, excretion, blood clotting, and neuromuscular functions. Suboptimal intakes of calcium may hinder normal growth and can lead to manifestations of osteoporosis in older people, especially postmenopausal women³⁹. The high

potassium and low sodium content of edible snails should be expected to exhibit physiological benefits. Low potassium consumption is often associated with a variety of physiological disorders including respiratory, renal problems and hypertension. Further, edible snails are found to be rich in micro-minerals like iron and zinc. The most vulnerable sections of the population suffering from iron deficiency are infants of the weaning stage, young children, and women of childbearing age. More than 1.2 billion people are suffering from iron deficiency anaemia and iron deficiency in the absence of anaemia is even more frequent³⁸. Iron deficiency anaemia is often caused by the low dietary intake of iron and reduced intake of bio-available iron, particularly in developing countries⁴⁰. Zinc is an essential component of a large number of enzymes (more than 300). Moreover, it has anti-oxidative stress and anti-inflammatory functions⁴¹. Copper, together with zinc, are additional structural components of the antioxidant enzyme superoxide dismutase. Considerable amounts of magnesium, phosphorus, and manganese are also present in edible snails. However, to study the bioavailability of the individual minerals from the snail meat remains a task of the future.

Safety issues

Although edible snails are nutritionally comparable, if not superior to conventional foods of animal origin, proper care has to be taken in relation to safety issues of using snails as food. Accumulation of heavy metals in freshwater snails is one important issue. Lau *et al.*, reported that two aquatic snail *Brotia costula* and *Clithon* sp. tissues contained arsenic contents that were higher than the permissible limit for human consumption⁴². In another experiment on *Pomacea canaliculata* exposed to contaminate




Fig. 6 — Content of macro-minerals a) calcium, magnesium, sodium, potassium, and phosphorus (mg/100 g) and content of micro-minerals; and b) iron, zinc, copper and manganese content (mg/100 g) of edible snail species. [Graphical representation was plotted based on Table 5].

Table 5 — Mineral content of edible snails

Scientific name	Mineral content (mg/100 g)									Reference
	Ca	Mg	Na	K	P	Fe	Zn	Cu	Mn	
<i>Limicolaria aurora</i>	401(DM)	771	178	533	636	ND	259	--	--	51
<i>Archachatina marginata</i>	212.38 (FM)	46.33	196.24	44.18	0.64	9.31	1.44	0.64	0.41	52
<i>Archatina</i> sp.	188.96	37.79	181.87	49.88	215.48	4.95	1.87	0.62	0.38	
<i>Limicolaria</i> sp.	22.24	18.91	53.83	72.82	183.35	4.66	1.23	0.16	0.59	
<i>Helix aspersa</i> (Foot)	200 (FM)	52.7	--	--	73	1	1	3.1	--	92
	1620 (DM)	425			590	8.4	8	25.2		
<i>Helix aspersa</i> (Viscera)	328	183	--	--	536	6.4	46.3	1.7	--	
	1120	625			1830	22	158	6		
<i>Helix aspersa maxima</i> (Foot)	183	76	--	--	83	1.4	0.8	2.1	--	
	1390	575			630	11.2	6.4	16.4		
<i>Helix aspersa maxima</i> (Viscera)	356	182	--	--	473	10.2	43.1	1.2	--	
	1370	700			1820	39.2	166	4.8		
<i>Helix lucorum</i> (Foot)	279	60.8	--	--	66.8	0.9	1.3	3.4	--	
	1840	400			440	6.0	8.8	22.8		
<i>Helixlucorum</i> (Viscera)	260	93.7	--	--	340	18	35.5	1.7	--	
	1040	375			1360	72	142	6.8		
<i>Helix pomatia</i> (Foot)	613	50.2	--	--	140	1.9	1.1	4	--	
	4580	375			1049	14.4	8.8	30		
<i>Helix pomatia</i> (Viscera)	328	134	--	--	487	7.6	60	1	--	
	1340	550			1990	31.2	248	4.4		
<i>Lymnaea stagnalis</i> (Meat)	118.8 (DM)	56.7	82.8	399	--	0.54	--	0.26	--	55
<i>Archachatina marginata</i> (ovum) Pfeiffer	201.08(FM)	45.59	50.80	192.78	123.43	8.69	1.54	ND	0.39	56
<i>Archachatina marginata</i> (saturalis) Philippi	207.53	45.34	52.93	209.95	123.23	9.41	1.69	ND	0.38	
<i>Achatina achatina</i>	204.63	46.15	60.94	193.74	131.38	9.43	1.76	ND	0.39	
<i>Limicolaria</i> sp.	208.75	45.99	65.10	197.57	153.89	9.46	1.51	ND	0.38	
<i>Archachatina marginata</i> (ovum)	180.71(DM)	29.2	--	60.79	59.16	6.79	1.19	--	--	57
<i>Archachatina marginata</i> (saturalis)	179.09	28.52	--	70	51.49	8.25	1.18	--	--	
<i>Limicolaria</i> sp.	172.79	27.33	--	64.52	59.79	9	1.14	--	--	
<i>Lanistes varicus</i>	152.7	30.56	--	69.39	62.52	10.1	1.3	--	--	
<i>Nucella lapillus</i>	181.49	31.73	--	72.4	60.19	11.09	1.32	--	--	
<i>Limicolaria</i> sp.	36.2 (FM)	5.28	--	--	8.98	0.72	--	0.29	--	58
<i>Achatina fulica</i>	66.3	15.13	--	--	14.79	1.3	--	0.58	--	
<i>Achatina achatina</i>	106.3	19.28	--	--	19.01	1.88	--	0.77	--	
<i>Archachatina marginata</i>	126.4	25.01	--	--	22.91	2.29	--	1.03	--	
<i>Achatina fulica</i>	186.35	--	28.97	64.15	62.73	1.51	--	--	--	59
	186.32	--	28.94	64.15	62.76	1.49	--	--	--	
<i>Limicolaria</i> sp.	179.65	--	24.78	56.72	58.34	1.36	--	--	--	
	179.64	--	24.83	56.76	58.32	1.36	--	--	--	
<i>Helix pomatia</i>	13.51	--	13.63	31.59	31.17	1.17	--	--	--	
	132.48	--	13.65	31.57	31.14	1.14	--	--	--	
<i>Hexaplex trunculus</i>	674.4 (FM)	178.7	196.1	224.8	95.2	8.1	112.8	3.1	0.68	60
<i>Helix aspersa</i>	135.70 (FM)	17.05	91.95	105.40	96.72	0.52	--	--	--	79
<i>Achatina achatina</i>	585.5 (DM)	--	--	331.8	269.2	9.8	39	3.3	--	63
<i>Cookia sulcata</i> (Small)	140 (DM)	330	400	300	110	5.81	3.22	0.64	--	84
<i>Cookia sulcata</i> (Large)	70	310	410	280	90	7.41	3.82	0.78	--	
<i>Pomacea canaliculata</i>	5161.2 (DM)	56.9	93.4	364.4	550.5	45.5	10.1	7.1	2	82
<i>Lunella torquata</i>	239 (FM)	69	301	305	153	3.24	1.4	0.11	0.05	85
<i>Lunella undulata</i>	44	65	270	333	164	4.11	1.52	0.06	0.05	
<i>Turbo militaris</i>	61	77	400	273	122	1.93	1.22	0.22	0.05	

(Contd.)

Table 5 — Mineral content of edible snails (*Contd.*)

Scientific name	Mineral content (mg/100 g)									Reference
	Ca	Mg	Na	K	P	Fe	Zn	Cu	Mn	
<i>Limicolaria flammea</i>	165.45	13.57	66.87	132.45	336.53	4.24	2.92	2.53	6.47	64
<i>Archachatina marginata</i> (Adult)	42.19	59.23	44.75	92.34	295.64	9.53	--	--	--	94
<i>Archachatina marginata</i> (Grower)	31.64	60.04	39.57	77.55	286.65	7.37	--	--	--	
<i>Archachatina marginata</i> (Snaillet)	26.46	238.8	31.26	69.24	274.5	5.25	--	--	--	

ND = Not detected
DM= Dry matter basis
FM= Fresh weight basis

sediment for two months it was revealed that higher than desired amounts of metals like iron, manganese, copper, and zinc had accumulated in the digestive tracts and glands⁴³. A similar trend was reported for *Bellamya* sp.⁴⁴. Onuoha *et al.*, stated that snails of the species *Archachatina marginata* collected from the crude oil-producing regions in Rivers State, Nigeria were not safe for human consumption, because of the carcinogenic risk posed by the high amounts of nickel in them⁴⁵. Besides, edible snails can host a number of parasites^{28,46}. In a study by Tesana *et al.*, from Northeast Thailand, the edible snail species *Pila polita*, *P. pesmei*, and *Hemiplecta distincta* all contained the nematode worm *Angiostrongylus scantonensis*, which can be transmitted to human hosts and cause Angiostrongyliasis⁴⁷. *Pomacea canaliculata* and *Achatina fulica* were found to be the most likely intermediate hosts of *Angiostrongylus cantonensis* in China⁴⁸. Among some other occasionally eaten snails, the species, *Limicolaria aurora*, *L. flammea*, and *Limicolariaopsis* sp. were found to be contaminated with mesocercariae of the trematode *Alaria*, while *L. flammea* was furthermore found to contain cercariae of the liver fluke *Dicrocoelium dendriticum*⁴⁶. Therefore, in order to avoid these hazards, maintaining sanitation of the snail farm is necessary and proper processing of edible snails is also important⁴⁹.

Conclusion

Although snail farming is not a novel entrant in agriculture, it is far from receiving proper appreciation. Various species of snails are used for food and/or medicinal purposes, but their roles are primarily confined to within the limitation of a few users in a small number of countries. Often dietary transitions due to urbanization, availability of western

foods, climate change, etc. are the major factors for the decline in the interest and the use of traditional foods including snails. To halt the erosion of the traditional food wisdom and to support the safe supply of snail meat, the establishment of helicicultural facilities is therefore, a necessity. An increase in snail meat consumption can mitigate nutrient deficiency and also enhance the economic condition of certain local areas in which suitable snails can be farmed and traded.

Acknowledgement

The study was partially supported by the BSRP grant through the National Research Foundation of Korea (NRF), Ministry of Education, grant number NRF-2018R1A6A1A03024862 received by Prof. Chuleui Jung.

Conflict of interest

The authors declare no conflict of interest.

Reference

- 1 von Grebmer K, Bernstein J, Resnick D, Wiemers M, Reiner L, *et al.*, 2022 Global hunger index: Food systems transformation and local governance. Bonn: Welthungerhilfe; and Dublin: Concern Worldwide, 2022, 1-54. Available from: <https://www.globalhungerindex.org/pdf/en/2022.pdf>, accessed on 3rd November 2022.
- 2 FAO, ICN 2 Second international conference on nutrition better nutrition better lives (FAO, Rome), 2014. Available from: <http://www.fao.org/about/meetings/icn2/toolkit/key-messages/en/>, accessed on 13th March 2021
- 3 Meyer-Rochow V B, Can insects help to ease the problem of world food shortage?, *Search*, 1975, 6(7), 261-262.
- 4 van Huis A, Itterbeeck J V, Klunder H, Mertens E, Halloran A, *et al.*, Edible insects: Future prospects for food and feed security food and agriculture organization of the united nations Rome, 2013 FAO Forestry Paper 171 (FAO, Rome), 2013, 1-187. Available from: <http://www.fao.org/3/i3253e/i3253e00.pdf>, accessed on 13th March 2021.
- 5 Hashemian M, Ahmadzadeh H, Hosseini M, Lyon S and Pourianfar H R, Production of microalgae-derived high-protein biomass to enhance food for animal feedstock and

- human consumption, in *Advanced Bioprocessing for Alternative Fuels, Biobased Chemicals, and Bioproducts. Technologies and Approaches for Scale-Up and Commercialization*, edited by M Hosseinni (Woodhead Publishing Series in Energy), 2019, 393-405.
- 6 Lubell D, Prehistoric edible land snails in the circum-Mediterranean: The archaeological evidence. *Petits animaux et sociétés humaines. Du complément alimentaire aux ressources utilitaires*. Éditions APDCA, Antibes, 2004, 77-98. Available from: <http://www.arts.uwaterloo.ca/~dlubell/Antibes.pdf>, accessed on 21st January 2021.
 - 7 Caetano D, Miranda A, Lopes S, Paiva J, Rodrigues A, *et al.*, Nutritional and toxicity profiles of two species of land snails *Theba pisana* and *Otala lactaea*, from Morocco, *J Food Compos Anal*, 2021, **100**, 103893.
 - 8 FAOSTAT, Food and agriculture data 2021. Rome: Food and agriculture organisation of United Nations (FAO, Rome), 2021. Available from: <http://www.fao.org/faostat/en/#data/QL>, accessed on 13th March 2021.
 - 9 Touchstone snails, Professionally specialized in snail business. Web based. Available from: www.touchstonesnailfranchise.com, accessed on 13th March 2021.
 - 10 Ghosh S, Meyer-Rochow V B and Jung C, Importance of neglected traditional food to ensure health and well-being, *Nutr Food Sci Int J*, 2018, **8**, 555729.
 - 11 Halwart M, The golden apple snail *Pomacea canaliculata* in Asian rice farming systems: Present impact and future threat, *Int J Pest Manag*, 1994, **40**, 199-206.
 - 12 Catalma M T E, Capil D T, Antalán R A, Serra A B, Barroga A J, *et al.*, Golden snail (*Pomacea* sp.) use in animal feeds, *Int Rice Res Newsllett*, 1991, **16**, 26.
 - 13 June L, Ulep L and Buenafe M M, Performance of broilers fed with snail (*Pomacea canaliculata*) meal as substitute to fish meal or meat and bone meal, *Tropicultura*, 1991, **9**, 58-60.
 - 14 Baghele M, Mishra S, Meyer-Rochow V B, Jung C and Ghosh S, Utilization of snails as food and therapeutic agents by Baiga tribals of Baihar tehsil of Balaghat district, Madhya Pradesh, India, *Int J Ind Entomol*, 2021, **43**, 78-84.
 - 15 Martin G H G, Bushmeat in Nigeria as a natural resource with environmental implications, *Environ Conserv*, 1983, **10**, 125-132.
 - 16 Cobbinah J R, *Snail Farming in West Africa, a Practical Guide* (Food and agriculture organization, Rome), 1983, 1-49. Available from: <http://online.fliphtml5.com/qjfd/kvoy/#p=4>, accessed on 13th March 2021.
 - 17 Cobbinah J R, Vink A and Onwuka B, *Snail farming: Production, processing and marketing*, (Agrodok 47), 2008.
 - 18 Adeyeye S A O, Bolaji O T, Abegunde T A and Adesina T O, Processing and utilization of snail meat in alleviating protein malnutrition in Africa: A review, *Nutr Food Sci*, 2020, **50**, 1085-1097.
 - 19 Ghosh S, Meyer-Rochow V B and Jung C, Farming the edible aquatic snail *Pomacea canaliculata* as a mini-livestock, *Fishes*, 2022, **7**, 6.
 - 20 Forte A, Zucaro A, Vico G D and Fierro A, Carbon footprint of heliciculture: A case study from an Italian experimental farm, *Agric Syst*, 2016, **142**, 99-111.
 - 21 Allen M J, *Molluscs in archaeology methods, approaches and applications*, (Oxbow Books), 2017.
 - 22 Meyer-Rochow V B, Therapeutic arthropods and other, largely terrestrial, folk-medicinally important invertebrates: A comparative survey and review, *J Ethnobiol Ethnomed*, 2017, **13**, 9.
 - 23 Prabhakar A K and Roy S P, Ethno-medicinal use of some shell fishes by people of Kosi river basin of North-Bihar, India, *Stud Ethno-Med*, 2009, **3**(1), 1-4.
 - 24 Bagde N and Jain S, Traditional and ethno zoological practices by tribes and rurals of Chindwara district of Madhya Pradesh, India, *World J Pharm Med Res*, 2017, **3**, 263-268.
 - 25 Kashyap V R, Ethno-zoological study in Gondwana Vindhyan region (Madhya Pradesh) India, *Int J Zool Stud*, 2017, **2**, 66-67.
 - 26 Pyron M and Brown K M, Introduction to mollusca and the class Gastropoda, in *Throp and Covich's Freshwater Invertebrates: Ecology and General Biology*, VI edn, 2015, 381-421.
 - 27 Madsen H and Hung N M, An overview of freshwater snails in Asia with main focus on Vietnam, *Acta Tropica*, 2014, **140**, 105-117.
 - 28 Meyer-Rochow V B, Snails (Terrestrial and Freshwater) as Human Food, in *Encyclopaedia of Food Security and Sustainability*, vol. 1, edited by P Ferranti, E M Berry & J R Anderson (Elsevier), 2019, 376-378.
 - 29 Mottet A, de Haan C, Falcucci A, Tempio G, Opio C, *et al.*, Livestock: On our plates or eating at our table? A new analysis of the feed/food debate, *Glob Food Sec*, 2017, **14**, 1-8.
 - 30 Murphy S P and Allen L H, Nutritional importance of animal sources foods, *J Nutr*, 2003, **133**, 3932S-3935S.
 - 31 Henchion M, Hayes M, Mullen A M, Fenelon M and Tiwari B, Future protein supply and demand: Strategies and factors influencing a sustainable equilibrium, *Foods*, 2017, **6**(7), 53.
 - 32 Kim S W, Less J F, Wang L, Yan T, Kiron V, *et al.*, Meeting global feed protein demand: Challenge, opportunity, and strategy, *Annu Rev Anim Biosci*, 2019, **7**, 17.1-17.23.
 - 33 RööS E, Sundberg C, Tidåker P, Strid I and Hansson P A, Can carbon footprint serve as an indicator of the environmental impact of meat production?, *Ecol Indic*, 2013, **24**, 573-581.
 - 34 Oonincx D G A B and de Boer I J M, Environmental impact of the production of mealworms as a protein source for humans-a life cycle assessment, *PLoS One*, 2012, **7**, 1-5.
 - 35 Grundy S M, Influence of stearic acid on cholesterol metabolism relative to other long-chain fatty acids, *Am J Nutr*, 1994, **60**, 986S-990S.
 - 36 Schneider C L, Cowles R L, Stuefer-Powell C L and Carr T P, Dietary stearic acid reduces cholesterol absorption and increases endogenous cholesterol excretion in hamsters fed cereal-based diets, *J Nutr*, 2000, **130**, 1232-1238.
 - 37 Mensink R P, Effects of stearic acid on plasma lipid and lipoproteins in humans, *Lipids*, 2005, **40**, 1201-1205.
 - 38 Van Rooijen M A and Mensink R P, Palmitic acid versus stearic acid: Effects of interesterification and intakes on cardiometabolic risk markers- A systematic review, *Nutrients*, 2020, **12**(3), 615.
 - 39 Liu C, Kuang X, Li K, Guo X, Deng Q, *et al.*, Effects of combined calcium and vitamin D supplementation on osteoporosis in postmenopausal women: A systematic review

- and meta-analysis of randomized controlled trials, *Food Funct*, 2020, **11**, 10817-10827.
- 40 Camaschella C, Iron metabolism and its disorders, *Blood*, 2019, **133**, 30-39.
- 41 Choi S, Liu X and Pan Z, Zinc deficiency and cellular oxidative stress: Prognostic implications in cardiovascular diseases, *Acta Pharmacol Sin*, 2018, **39**, 1120-1132.
- 42 Lau S, Mohamed M, Yen A T C and Su'ut S, Accumulation of heavy metals in freshwater molluscs, *Sci Total Environ*, 1998, **214**, 113-121.
- 43 Dummee V, Kruatrachue M, Trinachartvanit W, Tanhan P, Pokethitiyook P, *et al.*, Bioaccumulation of heavy metals in water, sediments, aquatic plant and histopathological effects on the golden apple snail in Beung Boraphet reservoir, Thailand, *Ecotoxicol Environ Saf*, 2012, **86**, 204-212.
- 44 Bo L, Wang D, Li T, Li Y, Zhang G, *et al.*, Accumulation and risk assessment of heavy metals in water, sediments, and aquatic organisms in rural rivers in the Taihu Lake region, China, *Environ Sci Pollut Res*, 2015, **22**, 6721-6731.
- 45 Onuoha S C, Anelo P C and Nkpaa K W, Human health risk assessment of heavy metals in Snail (*Archachatina marginata*) from four contaminated regions in rivers state, Nigeria, *Am Chem Sci J*, 2016, **11**, 1-8.
- 46 Igbinosa I B, Isaac C, Adamu H O and Adeleke G, Parasites of edible land snails in Edo state, Nigeria, *Helminthologia*, 2016, **53**, 331-335.
- 47 Tesana S, Srisawangwong T, Sithithaworn P, Laha T and Andrews R, Prevalence and intensity of infection with third stage larvae of *Angiostrongylus cantonensis* in mollusks from Northeast Thailand, *Am J Trop Med Hyg*, 2009, **80**(6), 983-987.
- 48 Song L, Wang X, Yang Z, Lv Z and Wu Z, *Angiostrongylus cantonensis* in the vector snails *Pomacea canaliculata* and *Achatina fulica* in China: A meta-analysis, *Parasitol Res*, 2016, **115**, 913-923.
- 49 Ghosh S, Jung C and Meyer-Rochow V B, Snail farming: An Indian perspective of a potential tool for food security, *Ann Aquacult Res*, 2016, **3**(3), 1024.
- 50 Andrews W H, Wilson C R, Romero A and Poelma P L, The Moroccan food snail, *Helix aspersa*, as a source of *Salmonella*, *Appl Microbiol*, 1975, **29**, 328-330.
- 51 Udoh A P, Akpanyung E O and Igiran I E, Nutrients and anti-nutrients in small snails (*Limicolaria aurora*), *Food Chem*, 1995, **53**, 239-241.
- 52 Adeyeye E I, Waste yield, proximate and mineral composition of three different types of land snails found in Nigeria, *Int J Food Sci Nutr*, 1996, **47**, 111-116.
- 53 Tetey E C T, Osei-Yaw A and Hodari-Okae M, Studies on the quality of traditionally smoked-dry snail meat (*Achatina achatina*) in Ghana, *Ghana J Agric Sci*, 1997, **30**, 145-150.
- 54 Adeyeye E I and Afolabi E O, Amino acid composition of three different types of land snails consumed in Nigeria, *Food Chem*, 2004, **85**, 535-539.
- 55 Yusuf A A and Oseni O A, Nutritional value and functional properties of pond snail (*Lymnaea stagnalis*), *Proc Int Conf Sci Nation Dev*, 2004, 90-93.
- 56 Fagbuaro O, Oso J A, Edward J B and Ogunleye R F, Nutritional status of four species of giant land snails in Nigeria, *J Zhejiang Univ Sci B*, 2006, **7**, 686-689.
- 57 Eneji C A, Ogogo A U, Emmanuel-Ikpeme C A and Okon O E, Nutritional assessment of some Nigerian land and water snail species, *Ethiop J Environ Stud Manag*, 2008, **1**, 56-60.
- 58 Babalola O O and Akinsoyinu A O, Proximate and mineral profile of snail meat from different breeds of land snail in Nigeria, *Pak J Nutr*, 2009, **8**, 1842-1844.
- 59 Adegoke A A, Bukola A T C, Comfort I U, Olayinka A A and Amos K O, Snails as meat source: Epidemiological and nutritional perspectives, *J Microbiol Antimicrobial*, 2010, **2**, 1-5.
- 60 Zarai Z, Frikha F, Balti R, Miled N, Gargouri Y, *et al.*, Nutrient composition of the marine snail (*Hexaplex trunculus*) from the Tunisian Mediterranean coasts, *J Sci Food Agric*, 2011, **91**, 1265-1270.
- 61 Malik A A, Aremu A, Bayode G B and Ibrahim B A, A nutritional and organoleptic assessment of the meat of the giant African land snail (*Archachatina marginata* Swainson) compared to the meat of other livestock, *Livest Res Rural Dev*, 2011, **23**(3), 60.
- 62 Kalio G A and Etela I, Nutritional and sensory profiling of the African giant land snails fed commercial –Type and leaf-based diets in a rain-forest ecology, *Afr J Food Agric Nutr Dev*, 2011, **11**, 5254-5265.
- 63 Engmann F N, Afoakwah N A, Darko P O and Sefah W, Proximate and mineral composition of snail (*Achatina achatina*) meat; any nutritional justification for acclaimed health benefits?, *J Basic Appl Sci Res*, 2013, **3**, 8-15.
- 64 Envin A J B, Ekissi G S E, Sea B T and Kouame P L, Biochemical and nutritional composition of garden snail (*Limicolaria flammea*) flesh consumed in Côte d'Ivoire, *J Basic Appl Re*, 2018, **4**, 63-70.
- 65 Bamidele J A, Ademolu K O, Idowu A B, Aladesida A A and Oladele A O, Biochemical and nutritional composition of giant African land snail (*Archachatina marginata*) from southwest Nigeria, *Trop Agric Sci*, 2018, **41**, 129-138.
- 66 Babatunde T O, Kehinde A S and Babatunde O O, Marketing analysis of snail in selected market in Ibadan metropolis, Oyo state, Nigeria, *J Res For Wildl Environ*, 2019, **11**, 106-113.
- 67 Meffowoet C P, Kouam K M, Kana J R and Tchakounte F M, Infestation rate of African giant snails (*Achatina fulica* and *Achatina marginata*) by parasites during the rainy season in three localities of Cameroon, *J Vet Med Res*, 2020, **7**, 1187.
- 68 Liat L B, Fong Y L, Krishnansamy M, Ramachandran P and Mansor S, Freshwater snail consumption and angiostrongyliasis in Malaya, *Trop Geogr Med*, 1978, **30**, 241-246.
- 69 Park I W and Kim C K, Fatty acid composition of *Achatina fucica* Bowdich and *Ampullarius insularis*, *J Korean Soc Food Nutr*, 1992, **21**(1), 36-42.
- 70 Lee M K, Moon J H and Ryu H S, Nutrient composition and protein quality of giant snail products, *J Korean Soc Food Nutr*, 1994, **23**, 453-458.
- 71 Saha B K, *Ecology and Bio-Economics of the Freshwater Edible Snails of Bangladesh*. Ph. D. Thesis, Rajshahi University, Rajshahi, 1998.
- 72 Nargis A, Talukdar D, Pramanik S H A and Hasan M R, Nutritional value and physico-chemical characteristics of apple snail *Pila globose* (Swainson) and *Lymnaea luteola* Lamark, *Bangladesh J Sci Ind Res*, 2011, **46**, 539-542.

- 73 Yusa Y and Wada T, Impact of the introduction of apple snails and their control in Japan, *Naga*, 1999, **22**(3), 9-13.
- 74 Özogul Y, Özogul F and Olgunnoglu A I, Fatty acid profile and mineral content of the wild snail (*Helix pomatia*) from the region of the south of the Turkey, *Eur Food Res Technol*, 2005, **221**, 547-549.
- 75 Lee S M and Lim T J, Effects of dietary protein and energy levels on growth and lipid consumption of juvenile snail (*Semisulcospira gottschei*), *J Shellfish Res*, 2005, **24**, 99-102.
- 76 Lv S, Zhang Y, Steinmann P and Zhou X N, Emerging angiostrongyliasis in mainland China, *Emerg Infect Dis*, 2008, **14**, 161-164.
- 77 Borkakati R N, Gogoi R and Borah B K, Snail: From present perspective to the history of Assam, *Asian Agri-Hist*, 2009, **13**(3), 227-234.
- 78 Lee M H, Kim Y K, Moon H S, Kim Y A, Yoon N Y, *et al.*, Antioxidant activities of five melania snails of the genus *Semisulcospira* in Korea, *Kor J Fish Aquat Sci*, 2010, **43**(3), 188-194.
- 79 Çağıltay F, Erkan N, Tosun D and Selçuk A, Amino acid, fatty acid, vitamin and mineral contents of the edible garden snail (*Helix aspersa*), *J Fish Sci*, 2011, **5**, 354-363.
- 80 Köhler F, Seddon M, Bogan A E, Tu D V, Sri-Aroon P, *et al.*, The status and distribution of freshwater molluscs of the Inolo-Burma region, in *The status and distribution of freshwater molluscs of the Indo-Burma region*, edited by D J Allen, K G Smith & W R T Darwall, (IUCN, Cambridge), 2012.
- 81 Debnath C, Sahoo L, Haldar A, Datta M, Yadav G S, *et al.*, Proximate and mineral composition of freshwater snails of Tripura, North-East India, *Fish Technol*, 2016, **53**, 307-312.
- 82 Ghosh S, Jung C and Meyer-Rochow V B, Snail as mini-livestock: nutritional potential of farmed *Pomacea canaliculata* (Ampullariidae), *Agric Nat Resour*, 2017, **51**, 504-511.
- 83 Ghosh-Jerath S, Kapoor R, Barman S, Singh G, Singh A, *et al.*, Traditional food environment and factors affecting indigenous food consumption in Munda tribal community of Jharkhand, India, *Front J Nutr*, 2021, **7**, 600470.
- 84 Mason S L, Shi J, Bekhit A E D and Gooneratne R, Nutritional and toxicological studies of New Zealand *Cookia sulcata*, *J Food Comp Anal*, 2014, **36**, 79-84.
- 85 Lah R A, Smith J, Savins D, Dowell A, Bucher D, *et al.*, Investigation of nutritional properties of three species of marine turban snails for human consumption, *Food Sci Nutr*, 2017, **5**, 14-30.
- 86 Elmslie L J, Snail collection and small-scale production in Africa and Europe, in *Ecological Implications of Mini livestock: Potential of Insects, Rodents, Frogs and Snails*, edited by M G Paoletti (Scientific Publishers Inc., Enfield), 2005, 93-121.
- 87 Ikauniece D, Jemeljanovs A, Sterna V and Strazdina V, Evaluation of nutrition value of Roman snail's (*Helix pomatia*) meat obtained in Latvia, in *Proc 9th Baltic Conference on Food Science and Technology "Food for Consumer Well-Being" Foodbalt*, 2014, 28-31.
- 88 Milinsk M C, Padre Rd G, Hayashi C, de Souza N E and Matsushita M, Influence of diets enriched with different vegetable oils on the fatty acid profiles of snail *Helix aspersa maxima*, *Food Chem*, 2003, **82**(4), 553-558.
- 89 Milinsk M C, Padre Rd G, Hayashi C, de Oliveira C C, Visentainer J V, *et al.*, Effects of feed protein and lipid contents on fatty acid profile of snail (*Helix aspersa maxima*) meat, *J Food Compos Anal*, 2006, **19**, 212-216.
- 90 Watt B K and Merrill A L, *Composition of foods. Agriculture Handbook, 8* (US Department of Agriculture, Washington DC), 1975.
- 91 Claeys E and Demeyer D, Nutritional quality of snails (*Helix aspersa*), in *Proc II-32nd European Meeting of Meat Research Workers* (Ghent), 1986, 415-418.
- 92 Gomot A, Biochemical composition of *Helix* snails: Influence of genetic and physiological factors, *J Molluscan Stud*, 1998, **64**, 173-181.
- 93 Baby R L, Hasan I, Kabir K A and Naser M N, Nutrient analysis of some commercially important molluscs of Bangladesh, *J Sci Res*, 2010, **2**, 390-396.
- 94 Kehinde A S, Babatunde T O and Kehinde O J, Lifecycle evaluation of the nutritional benefits and biosafety of snail (*Archachatina marginata*), *J Exp Agric Int*, 2019, **40**, 1-7.
- 95 Ekin I, Başhan M and Şeşen R, Possible seasonal variation of the fatty acid composition from *Melanopsis praemorsa* (L., 1758) (Gastropoda: Prosobranchia), from southeast Anatolia, Turkey, *Turk J Biol*, 2011, **35**, 203-213.
- 96 EkinI, A comparative study on fatty acid content of main organs and lipid classes of land snails *Assyriella escheriana* and *Assyriella guttata* distributed in Southeastern Anatolia, *Ital J Food Sci*, 2015, **27**, 1-7.
- 97 Ekin I and Şeşen R, Investigation of the fatty acid contents of edible snails *Helix lucorum*, *Eobania vermiculata* and non-edible slug *Limax flavus*, *Rec Nat Prod*, 2017, **11**, 562-567.
- 98 Szkucik K, Ziomek M, Paszkiewicz W, Drozd Ł, Gondek M, *et al.*, Fatty acid profile in fat obtained from edible part of land snails harvested in Poland, *J Vet Res*, 2018, **62**(4), 519-526.
- 99 Galluzzo F G, Cammilleri G, Ulrici A, Calvini R, Pulvirenti A, *et al.*, Land snails as a valuable source of fatty acids: A multivariate statistical approach, *Foods*, 2019, **8**(12), 676.