Indian Journal of Traditional Knowledge
Vol 25(10), October 2026, pp 975-983
DOI: 10.56042/ijtk.v25i10.23689

Ameliorative potential of *Katuka Musali Kashay* (polyherbal decoction of *Picrorhiza kurroa* and *Asparagus adscendens*) in methotrexate-induced hepatotoxicity in Wistar rats

Tanvi Harvadan Umrigar[a,*] & Sunila H Deo[b]

[a]Department of Agadtantra, P P Savani Ayurvedic College and Hospital, P P Savani University, Surat, Gujarat 394 125, India
[b]Department of Agadtantra, Dr D Y Patil College of Ayurved and Research Centre, Pimpri, Pune, Maharashtra 411 018, India
[*]E-mail: tanviumrigar1@gmail.com

Received 11 September 2025; revised 05 September 2026; accepted 12 October 2026

Supplementary Data




Fig. S1 — Effect of *Katuka Musali Kashay* on body weight gain (g) in MTX-induced hepatotoxicity. Day 7–0: weight gain from Day 0 to Day 7; Day 14–7: from Day 7 to Day 14; Day 14–0: from Day 0 to Day 14. Two-way ANOVA followed by Bonferroni post hoc test




Fig. S2 — Serum SGOT (IU/L)




Fig. S3 — Serum SGPT (IU/L)




Fig. S4 — Serum bilirubin (mg/dL)




Fig. S5 — Serum total protein (g/dL)




Fig. S6 — Serum ALP (IU/L)




Fig. S7 — Liver MDA (µM/mg tissue). One-way ANOVA followed by Tukey's post hoc test




Fig. S8 — Liver GSH (µM/mg tissue). One-way ANOVA followed by Tukey's post hoc test