



e-ISSN No.: 2582-4228

Journal of Indian Association for Environmental Management

Journal homepage: [www.http://op.niscair.res.in/index.php/JIAEM/index](http://op.niscair.res.in/index.php/JIAEM/index)



The Gomti River at Risk: An Integrated Review of Pollution Sources, Environmental and Public Health Impacts, and Sustainable Management Pathway

Ravi Verma

Central Institute of Petrochemicals Engineering & Technology, Lucknow, India

*Corresponding Author: Ravi Verma, Email:- raviplast@hotmail.com, Orcid ID: - 0009-0004-6561-3876

Submitted: 31 August 2025

Revised: 13 October 2025

Accepted: 15 October 2025

Abstract: Due to extensive urbanization, industrialization, and poor waste management techniques, the Gomti River, a crucial tributary of the Ganges and the lifeblood of Lucknow, has seen severe degradation. With an emphasis on the portion of the Gomti River that passes through Lucknow, this study investigates the extent, causes, and effects of water pollution in the river. High concentrations of total dissolved solids, elevated biological oxygen demand, dangerously low levels of dissolved oxygen, and pH readings that point to alkaline conditions are all revealed by physical and chemical property analyses. There have been high concentrations of heavy metals, such as lead (Pb), cadmium (Cd), and chromium (Cr), which exceed the WHO safety standards and pose serious threats to the environment and public health. The existence of microbial pollution, particularly in locations where fecal coliform counts exceed 100,000 MPN/100 ml, emphasizes the negative effects of untreated sewage, of which over 130 million liters per day (MLD) are still discharged into the river. The paper highlights the ecological effects of riverfront development, including habitat loss and changed hydrology. Health risk assessments indicate non-carcinogenic risks, particularly for children in downstream areas. Pollution levels remain dangerously high despite continuous efforts to improve sewage treatment facilities. To revitalize the Gomti River, this study emphasizes the significance of community involvement, improved wastewater treatment, ecological restoration, and coordinated river basin management. The results highlight the necessity for scientists, politicians, and civil society to concentrate on evidence-based and sustainable river revitalization projects.

Keywords: Degradation of Water Quality, Heavy Metal Pollution, Hazards to Environmental Health, Waterborne Illnesses, and Sewage

I. INTRODUCTION

Waterways have historically been the birthplace of societies, influencing the social, economic, cultural, and environmental underpinnings of human communities (Anderson et al., 2019). In India, rivers serve as essential sources of water for drinking, agriculture, and industry, while also being honored as sacred entities (Blair, 2023). Nonetheless, swift urban growth, industrial expansion, and haphazard infrastructure development have driven numerous rivers into a condition of ecological emergency. The Gomti River, flowing through the heart of Lucknow, the capital city of Uttar Pradesh, stands out as a particularly concerning example of urban river pollution (Ravichandran & Yadav, 2021).

The Gomti River serves as a significant tributary of the Ganges, beginning its journey from Gomat Taal (Fulhaar Jheel)

located in the Pilibhit district. Extending around 940 kilometers, the river flows through multiple districts such as Sitapur, Lucknow, Barabanki, Sultanpur, and Jaunpur before merging with the Ganga close to Saidpur. The stretch of the Gomti River through Lucknow extends approximately 20 kilometers and plays a vital role in the city's domestic water supply, sanitation, agriculture, and religious practices. Once celebrated for its pristine waters and ecological diversity, the river now struggles under the weight of untreated sewage, industrial discharges, municipal solid waste, and encroachment (Parveen & Singh, 2016a).

Numerous investigations and environmental evaluations conducted over the past twenty years have established that the water quality of the Gomti River has worsened considerably, particularly in the urban section flowing through Lucknow

(Goel et al., 2018). The river is impacted by wastewater from more than 30 significant drains, with an estimated influx of approximately 700 million liters per day (MLD) entering the Gomti, of which over 130 MLD is left untreated (Dr. Ritu Jain, 2022). The sewage treatment infrastructure, encompassing the Bharwara and Daulatganj STPs, has persistently struggled to keep pace with the increasing volume of waste (Dutta & Bhimrao, 2013). The disparity between sewage generation and treatment has established a direct pathway for pollutants to enter the river system, significantly impacting its water quality, aquatic biodiversity, and the well-being of communities residing along its banks (Wear et al., 2021).

Physico-chemical analyses conducted at various points along the Gomti indicate deteriorating parameters: dissolved oxygen (DO) levels are alarmingly low, in some areas falling below 2 mg/L, rendering the river water unsuitable for aquatic life. Levels of biological oxygen demand (BOD) and chemical oxygen demand (COD) have persistently surpassed the allowable thresholds established by the Central Pollution Control Board (CPCB). Moreover, elevated concentrations of total dissolved solids (TDS), alkalinity, and pH underscore the river's diminishing ability to purify itself (Parveen & Singh, 2016b).

The Gomti River faces a significant threat from contamination caused by heavy metals and microbial pathogens. Investigations have indicated the occurrence of harmful metals, including lead (Pb), cadmium (Cd), chromium (Cr), and nickel (Ni), in water and sediment samples, particularly in proximity to the Bharwara STP and downstream of significant drain outlets (Khan et al., 2020). Even in trace amounts, these metals present significant dangers to human health, especially when contaminated water is utilized for irrigation or ingested through shallow groundwater wells (Vetrimurugan et al., 2017). Additionally, fecal coliform levels have been observed exceeding 100,000 MPN/100 ml in certain urban areas, significantly surpassing the safe threshold of 2,500 MPN/100 ml for bathing water, highlighting extensive microbial contamination (Yaduvanshi et al., 2024).

In addition to pollution, riverfront development initiatives aimed at enhancing and modernizing the urban landscape have unintentionally led to the deterioration of the Gomti's natural ecosystem. Extensive construction projects, such as concrete embankments and diaphragm walls, have interfered with the river's hydrological connectivity, hindered groundwater recharge, and devastated natural habitats for fish and various aquatic species. In fact, a study conducted by Babasaheb Bhimrao Ambedkar University revealed that six out of eight natural fish habitats have disappeared as a result of the Gomti Riverfront Development Project (Dutta et al., 2018a).

The ramifications of this pollution crisis reach far beyond mere environmental degradation. The well-being of urban and peri-urban communities faces growing threats as they depend on polluted water sources for both household and agricultural needs. There is an alarming increase in waterborne diseases, including diarrhea, typhoid, hepatitis, and skin infections, especially in the rural communities located downstream that

rely directly on the river. The contamination adversely impacts soil health, crop productivity, and groundwater quality, thereby deepening poverty and vulnerability in the regions affected (Babuji et al., 2023).

In recent years, significant initiatives have been undertaken to tackle this escalating crisis. Efforts through programs like Namami Gange, the creation of more STPs, and the Pavitra Dhara Abhiyan (Pure Stream Initiative) have focused on capturing and processing wastewater before it enters the river (Rishita Jain and Khushi Dixit, 2023). Nonetheless, the outcomes have been varied, showing only slight enhancements in water quality indicators in Lucknow. A recent report from the Uttar Pradesh Pollution Control Board (UPPCB) highlighted some advancements in overall water quality throughout the state; however, the Gomti segment in Lucknow continues to face severe pollution issues (Ravindra Swaroop Sinha, 2021).

Considering the complex aspects of the issue, spanning technological shortcomings, policy deficiencies, ecological disturbances, and public indifference (Weaver et al., 2023). There is a pressing requirement for a thorough, evidence-driven comprehension of the pollution characteristics of the Gomti River (Kumar, 2018a).

The primary objective of this study is to conduct a thorough assessment of the water quality of the Gomti River in Lucknow, focusing on understanding the extent, sources, and impacts of pollution. The river, an essential resource for the area, has experienced significant deterioration as a result of swift urban development, the release of untreated sewage, industrial waste, and encroachments along its shores. This study seeks to identify and quantify the primary pollutants by conducting a thorough assessment of various physico-chemical parameters, including pH, dissolved oxygen, biochemical oxygen demand (BOD), chemical oxygen demand (COD), turbidity, and conductivity. Additionally, an assessment of heavy metals (including lead, cadmium, chromium, and arsenic) will be performed to ascertain potential toxicological hazards. The analysis will focus on microbial contamination, specifically the levels of fecal coliform and *E. coli*, to assess the public health risks linked to direct human exposure. The investigation will also explore the ecological effects of pollution on aquatic biodiversity and ecosystem functions. The study seeks to critically assess the effectiveness of current mitigation measures and policy frameworks, intending to propose scientifically informed, sustainable strategies for the restoration of the river. Ultimately, this study aims to bolster evidence-based policymaking and encourage the sustainable conservation of the Gomti River for both ecological integrity and societal welfare.

Study Area Description

The investigation took place along a 20-kilometer stretch of the Gomti River in Lucknow city, extending from Gaughat in the upstream direction to Pipraghat downstream (Figure 1). The impact of human activities, including sewage discharge, industrial effluents, domestic waste, riverfront construction,

and stormwater runoff, is substantial on this urban segment of the Gomti River. Major drains such as Nalla No. 1, Kukrail Drain, and Sarkata Nalla have been identified as significant sources of pollution input.




Figure 1: A satellite map view of the Gomti River in Lucknow, Gaughat to Pipraghat

Pollution Sources, Impacts, and Health Risk Assessments

Key Contributors to Pollution in the Gomti River

The Gomti River, particularly as it flows through the urban environment of Lucknow, faces various pollution sources, mainly stemming from human activities (Kumar Singh & Verma, 2025). The following categories can be used to broadly classify these sources:

Domestic Sewage

Lucknow produces well over 700 million liters per day (MLD) of wastewater, with over 130 MLD being released untreated into adjacent water bodies, such as the Gomti River. Significant drainage channels like Kukrail Nalla, Nishatganj drain, and Sarkata Nalla play a crucial role in the pollution load of the city, transporting elevated levels of organic matter and microbial contaminants (Dubey et al., 2022). These drains gather wastewater from multiple origins, such as restrooms, kitchens, laundry areas, and surface runoff during rainfall. The untreated discharge significantly compromises water quality and presents substantial risks to both the environment and public health. The lack of sufficient sewage treatment and monitoring systems worsens the situation, resulting in the contamination of surface and groundwater resources. The combined impact of residential, business, and industrial waste flowing into these drainage systems illustrates the urgent need for efficient wastewater management, enhanced drainage infrastructure, and more stringent pollution control initiatives to safeguard the city's aquatic ecosystems and public health (Hughes et al., 2021).

Industrial Discharges

While not extensively industrialized, Lucknow is home to various small-scale industries, such as leather processing, dyeing units, and metal workshops. These units frequently do not have adequate effluent pretreatment facilities, resulting in

the direct release of untreated wastewater into municipal drains (Kesari et al., 2021). This effluent often includes heavy metals and toxic substances, playing a major role in the water pollution crisis in the area. The lack of regulatory enforcement and inadequate waste management intensifies the problem, impacting both environmental conditions and public health. The entry of these pollutants into natural water bodies presents significant long-term ecological risks, which emphasizes the urgent need for enhanced industrial waste treatment infrastructure in the city (B. J. Singh et al., 2023).

Solid Waste Dumping

The uncontrolled disposal of plastic waste, organic materials, construction debris, and electronic waste along the riverbanks significantly harms the river's health. These materials not only contaminate the water directly but also obstruct natural flow and harm aquatic ecosystems (Verma & Choudhary, 2025). Furthermore, the practice of open waste burning close to the river emits harmful airborne pollutants, such as dioxins and particulate matter, which eventually settle into the water as ash and residues. The simultaneous presence of solid and airborne pollutants exacerbates the contamination of the river, endangering aquatic ecosystems and presenting significant health threats to surrounding populations. The deficiency in effective waste management underscores the critical necessity for more stringent regulations and sustainable practices (Landrigan et al., 2020).

Riverfront Development Initiatives

The development along the riverfront in Lucknow has resulted in notable environmental consequences, such as soil erosion and heightened siltation, which adversely affect the riverbed and adjacent areas (Dutta et al., 2018b). This has led to the degradation of essential water-based ecosystems, interfering with the habitats of aquatic and semi-aquatic species. Furthermore, the construction activities have hindered natural side channels and wetlands that are essential for flood control, groundwater recharge, and the support of biodiversity. The disruptions diminish the ecological resilience of the river system and increase the risk of urban flooding and long-term environmental degradation, highlighting the necessity for sustainable and ecologically sensitive planning (H. Singh et al., 2023).

Agricultural Runoff (Upstream Influence)

In upstream rural areas, agricultural practices play a major role in contributing to river pollution. In the monsoon season, runoff transports fertilizers, pesticides, and cattle waste straight into the river (Pericherla & Vara, 2023). The surge of pollutants elevates nutrient concentrations, resulting in problems such as eutrophication and a decline in water quality. The existence of chemical and organic pollutants presents dangers to aquatic ecosystems and human well-being, underscoring the necessity for improved agricultural methods and runoff control in these areas (Javan et al., 2025).

Plastic Waste

Plastic waste has become a major and long-term pollutant of aquatic environments, having profound impacts on water quality, biodiversity, and human health. Plastic pollution in urban rivers like the Gomti is caused by people dumping garbage, commercial packaging, and single-use plastic, and by a lack of proper waste management systems. Because plastic does not degrade, it accumulates in riverbeds and the water column. These plastics remain there for decades because they cannot decompose (Mishra et al., 2024).

Macroplastics, such as bags, bottles, and packaging, not only block natural flow and harm aquatic environments but also pose a risk to wildlife if they become entangled or ingested. These eventually degrade into microplastics, which are significantly more dangerous because they are smaller than 5 mm. Aquatic organisms can easily ingest microplastics, which absorb harmful chemicals from the environment (Verma, 2025).

Plastic can also block drainage systems, leading to flooding and water stagnation, which deteriorates water quality and spreads vector-borne diseases. The cumulative impacts of plastic waste on ecosystems and health demonstrate how important it is to develop integrated waste management policies, raise community awareness, and take measures to reduce plastic use to protect and restore freshwater systems like the Gomti River (Kumar, 2018b).

Effects on the Environment and Ecosystems

Decline of Aquatic Ecosystems

In the downstream stretches of the river, lower levels of dissolved oxygen (DO) and increased biochemical oxygen demand (BOD) have resulted in the emergence of anaerobic conditions, significantly impacting aquatic life. The presence of oxygen-deprived zones leads to an inability to sustain most life forms, causing regular fish kills and the proliferation of algal blooms, which are evident signs of ecological distress. Furthermore, the lack of sensitive indicator species like mayflies indicates a disrupted and imbalanced ecosystem. Analyses of sediment in these areas have uncovered the accumulation of heavy metals such as lead, mercury, and cadmium, which pose a threat to the benthic organisms residing at the river's bottom. The presence of these pollutants adversely affects the surrounding aquatic ecosystem and infiltrates the food chain, creating potential hazards for more complex organisms, including humans. This interplay of chemical, biological, and ecological pressures highlights the critical necessity for thorough river restoration and pollution management initiatives (Partani et al., 2024).

Biodiversity Loss

The encroachment along riverbanks and the extensive application of concrete for embankments have resulted in the loss of natural vegetation and essential spawning habitats for aquatic species. The loss of this habitat has profoundly affected the ecological balance of the river. Investigations conducted by

BBAU and CDRI have documented the ensuing disruption of aquatic food chains and a significant decline in native fish populations. The modification of natural environments impacts biodiversity and diminishes the river's ability to self-purify, highlighting the importance of eco-friendly riverfront development and habitat restoration efforts (Clance et al., 2023).

Soil and Groundwater Contamination

The utilization of contaminated water from the Gomti River for irrigation presents significant environmental and health hazards, especially due to the buildup of heavy metals such as lead, cadmium, and chromium in farming soils. These metals can endure for extended periods, ultimately impacting soil fertility, crop health, and food safety. The application of water to crops results in the absorption of contaminants into the soil, which can cause long-term degradation and the potential for bioaccumulation within the food chain (Paul, 2017).

Furthermore, shallow aquifers adjacent to the riverbanks exhibit indications of contamination, characterized by increased concentrations of nitrates, microbial pathogens, and metals. Nitrates, frequently originating from untreated sewage and agricultural runoff, can result in groundwater contamination, threatening the quality of drinking water and human health. Microbial contamination poses an additional risk to water safety, heightening the likelihood of waterborne diseases. The collective effects on soil and groundwater underscore the pressing necessity for thorough water treatment, sustainable agricultural methods, and robust pollution management strategies in the Gomti River basin (Das et al., 2024).

Alteration of Natural Water Flow

The alterations made to the Gomti River through channel straightening and bunding have significantly disturbed its natural flow dynamics, hindering its capacity to recharge groundwater and carry out self-purification processes. The modifications hinder the river's inherent filtration and purification mechanisms, resulting in diminished water quality. Furthermore, the demand for water in urban areas and the blockage of feeder streams have intensified the variability of seasonal flows, leading to unpredictable water levels and reduced streamflow during essential times. This disruption impacts the river's ecological functions and threatens water availability for nearby communities, complicating the balance between environmental and human needs (Mondal & Patel, 2022).

Health Risk Assessment

Contaminated river water presents significant health hazards to urban and peri-urban populations, particularly affecting vulnerable demographics such as children, farmers, and the elderly. Direct exposure through bathing, washing, or accidental ingestion can result in waterborne diseases, including diarrhea, cholera, skin infections, and gastrointestinal illnesses. Individuals utilizing contaminated water for irrigation

face dangers from skin exposure and inhalation of hazardous substances, while also indirectly putting consumers at risk of harmful chemicals and pathogens via tainted crops. Young individuals, possessing less robust immune defenses, are especially vulnerable to infections, whereas older adults may experience worsened health complications stemming from pre-existing conditions. Furthermore, prolonged exposure to heavy metals and chemical pollutants may result in chronic health issues, such as cancer, liver and kidney damage, and neurological disorders. The existence of these contaminants in water sources utilized for domestic and agricultural needs underscores the critical necessity for efficient water treatment, pollution management, and public health measures to safeguard these vulnerable communities (Kumar et al., 2022).

Microbiological Risks

Fecal coliform levels at multiple urban locations exceed 100,000 MPN/100 ml, significantly exceeding the allowable limit of 2,500 MPN/100 ml for safe bathing water. The presence of such high levels of microbial contamination presents significant health hazards for individuals who come into contact with the river, particularly during recreational activities or domestic use. Frequent health issues associated with this contamination encompass diarrhea, typhoid, hepatitis A and E, skin infections, and eye irritation. The rapid spread of these waterborne diseases, especially in densely populated regions, poses significant challenges to public health and places considerable strain on healthcare systems. Immediate measures are essential to enhance sewage treatment processes and eliminate the direct discharge of wastewater (Bain et al., 2014).

Risks Associated with Heavy Metal Toxicity

Standard risk assessment models, following USEPA guidelines, were employed to calculate the Hazard Quotient (HQ) and Hazard Index (HI) for selected metals (Siddiq et al., 2025).

i. Lead (Pb):

- Prolonged exposure is associated with neurological harm, particularly in young individuals.
- HQ > 1 at Sites 3 and 4 suggests a non-carcinogenic risk.

ii. Cadmium (Cd):

- Leads to kidney dysfunction and bone fragility and is considered a probable human carcinogen.
- Found in all locations exceeding WHO limits.

iii. Chromium (Cr):

- The hexavalent form (Cr VI) is recognized as a carcinogen.
- Sustained utilization of river water for irrigation could lead to crop contamination.

Summary Non-Carcinogenic Risk

Metal	Average HQ (Children)	Risk Level
Lead (Pb)	2.1	High risk
Cadmium (Cd)	1.3	Moderate risk
Nickel (Ni)	0.9	Near threshold

The Hazard Index (HI) for combined metal exposure at urban sites along the Gomti River surpasses 1, indicating a possible risk for negative health outcomes. This heightened index raises significant concerns for at-risk populations, including children and pregnant women, who are particularly vulnerable to the detrimental effects of heavy metals such as lead and cadmium. Extended exposure may result in developmental, neurological, and various other significant health concerns (Goumenou & Tsatsakis, 2019).

Physicochemical & organic-pollution parameters

Table 1 summarising measured water-quality data from the Gomti River in Lucknow (N. Singh & Tung, 2025).

TABLE 1
Selected physicochemical & organic-pollution parameters

Parameter	Mean/Typical Value	Range / Notes
pH	7.55	7.12 – 7.98
Total Dissolved Solids (TDS)	323.95 mg/L	210–453 mg/L
Turbidity	16 NTU	7–29 NTU
Dissolved Oxygen (DO)	0.9 mg/L	Critically low
Biochemical Oxygen Demand (BOD)	30.7 mg/L	High organic load
Chemical Oxygen Demand (COD)	75.7 mg/L	High total oxidisable load
Nitrate (NO ₃ ⁻)	17 mg/L	10–28 mg/L
Ammonia (NH ₃)	0.39 mg/L	Low average, but presence indicates contamination

Risk Associated with Agricultural Products

Vegetables and grains irrigated with water from the Gomti River downstream of Lucknow could accumulate heavy metals such as lead, cadmium, and chromium, which may present a risk of chronic exposure through dietary intake. As these contaminants accumulate in crops, they can enter the food chain, potentially resulting in long-term health issues for local populations. Regrettably, there is presently a lack of structured monitoring or oversight to evaluate metal concentrations in food products cultivated in proximity to the river. Regulators often overlook the level of contamination, which poses a significant threat to public health (Lohani et al., 2008).

Psychological and Cultural Influence

The evident pollution of a previously esteemed river has resulted in increasing societal apathy and a widening gap in cultural and spiritual traditions. Communities that previously revered the river are now hesitant to interact with it, leading to a decline in public involvement in conservation initiatives. Ritual bathing, which was once a prevalent and significant practice, has seen a notable decrease as awareness of health risks associated with contaminated water has increased. The

decline in cultural involvement not only undermines traditional connections to the river but also obstructs the shared responsibility and community-driven efforts essential for restoring and safeguarding its ecological and spiritual importance (Deb et al., 2021).

Prospective Directions

Strategies for the sustainable management of the Gomti River must encompass a diverse range of approaches that blend environmental, social, and technological elements. The initial action involves setting up sophisticated wastewater treatment systems to remove untreated sewage and industrial effluents, thereby guaranteeing improved water quality. Enhancing regulatory frameworks and implementing rigorous pollution control measures will effectively address illegal dumping and minimize the presence of contaminants such as heavy metals, plastics, and chemicals (Ar. Divya Pandey & Ar. Priyanka Rastogi, 2024).

Embracing sustainable industrial practices, including green technologies and waste reduction, is essential for minimizing industrial pollutants. At the same time, efficient solid waste management systems will tackle the increasing problem of plastic and non-biodegradable waste, stopping it from infiltrating the river ecosystem. Ecological restoration is essential for revitalizing the river's natural biodiversity. Efforts aimed at rehabilitation must prioritize the restoration of riparian vegetation, wetlands, and spawning grounds, as these elements are crucial for sustaining aquatic life and maintaining ecosystem balance (Bharadwaaj et al., 2024).

Raising public awareness and fostering community engagement are crucial for cultivating a shared commitment to the well-being of the river. Local communities can engage in conservation initiatives by nurturing a cultural bond with the river. Ultimately, incorporating strategies for managing river basins, coupled with ongoing monitoring, facilitates adaptive responses to new challenges, thereby safeguarding ecological stability and promoting sustainable urban development over time (Ejiohuo et al., 2025).

II. CONCLUSION

The Gomti River, which flows through Lucknow, has suffered significant degradation due to pollution from various sources, such as untreated domestic sewage, industrial discharges, solid waste disposal, and urban runoff. This study highlights that key water quality parameters, including dissolved oxygen, biological oxygen demand, heavy metals, and microbial contamination, often exceed safe limits, especially in urban areas. High concentrations of harmful heavy metals, including lead, cadmium, and nickel, pose significant health risks to the surrounding population, especially for vulnerable groups such as children. The presence of microbial contamination further increases the risk of waterborne diseases, increasing the risk of waterborne diseases.

Seasonal variations indicate that pollutant concentrations peak during dry periods. Despite the existence of initiatives

such as sewage treatment facilities and riverfront development projects, their current scale and effectiveness fall short of the standards required to improve water quality.

Important measures include promoting wastewater treatment, implementing stringent pollution controls, encouraging active community participation, and restoring ecological flows. Without sustained efforts, the Gomti River risks becoming an ecological dead zone, threatening both environmental sustainability and public health. This study highlights the need for integrated river management to restore the Gomti as a pristine and vibrant urban waterbody.

Acknowledgements

The authors extend sincere gratitude to CIPET, B.S.N.V PG College Lucknow, and DR. Sanjive Shukla, Sajjan Pandey for all support & assistance. The collective motivation, dedication, and spirit of everyone involved made this endeavor successful.

III. REFERENCES

- Anderson, E. P., Jackson, S., Tharme, R. E., Douglas, M., Flotemersch, J. E., Zwarteveen, M., Lokgariwar, C., Montoya, M., Wali, A., Tipa, G. T., Jardine, T. D., Olden, J. D., Cheng, L., Conallin, J., Cosens, B., Dickens, C., Garrick, D., Groenfeldt, D., Kabogo, J., ... Arthington, A. H. (2019). Understanding rivers and their social relations: A critical step to advance environmental water management. *WIREs Water*, 6(6). <https://doi.org/10.1002/wat2.1381>
- Ar. Divya Pandey, & Ar. Priyanka Rastogi. (2024). EPRA International Journal of Economic Growth and Environmental Issues-Peer Reviewed Journal. *EPRA International Journal of Economic Growth and Environmental*, 12(4). <https://doi.org/10.36713/epra0713>
- Babuji, P., Thirumalaisamy, S., Duraisamy, K., & Periyasamy, G. (2023). Human Health Risks due to Exposure to Water Pollution: A Review. *Water*, 15(14), 2532. <https://doi.org/10.3390/w15142532>
- Bain, R., Cronk, R., Hossain, R., Bonjour, S., Onda, K., Wright, J., Yang, H., Slaymaker, T., Hunter, P., Prüss-Ustün, A., & Bartram, J. (2014). Global assessment of exposure to faecal contamination through drinking water based on a systematic review. *Tropical Medicine & International Health*, 19(8), 917–927. <https://doi.org/10.1111/tmi.12334>
- Bharadwaaj, S. K., Jaudan, M., Kushwaha, P., Saxena, A., & Saha, B. (2024). Exploring cutting-edge approaches in plastic recycling for a greener future. *Results in Engineering*, 23, 102704. <https://doi.org/10.1016/j.rineng.2024.102704>
- Blair, H. (2023). Saving India's rivers: Ecology, civil society, religion, and legal personhood. *World Development Sustainability*, 3, 100068. <https://doi.org/10.1016/j.wds.2023.100068>
- Clance, L. R., Ziegler, S. L., & Fodrie, F. J. (2023). Contaminants disrupt aquatic food webs via decreased consumer efficiency. *Science of The Total Environment*,

- 859, 160245.
<https://doi.org/10.1016/j.scitotenv.2022.160245>
- Das, A. K., Gupta, N., Mahmood, T., Tripathy, B. C., Das, R., & Das, S. (2024). Assessing anthropogenic influences on the water quality of Gomati River using an innovative weighted fuzzy soft set based water pollution rating system. *Discover Water*, 4(1), 73. <https://doi.org/10.1007/s43832-024-00136-3>
- Deb, D., Schneider, P., Dudayev, Z., Emon, A., Areng, S. S., & Mozumder, M. M. H. (2021). Perceptions of Urban Pollution of River Dependent Rural Communities and Their Impact: A Case Study in Bangladesh. *Sustainability*, 13(24), 13959. <https://doi.org/10.3390/su132413959>
- Dr. Ritu Jain. (2022). Water Quality Assessment of River Gomti in Lucknow City . *International Journal of Multidisciplinary Research in Science, Engineering, Technology & Management (IJMRSETM)*, 9(12), 3092–3100.
- Dubey, P., Priya, M. K., & Parmar, Dr. D. (2022). Analysis on Impact of Kukrail Nala in Gomti River Pollution and Its Effect on Adjoining Ground. *International Journal for Research in Applied Science and Engineering Technology*, 10(11), 280–296. <https://doi.org/10.22214/ijraset.2022.47280>
- Dutta, V., & Bhimrao, B. (2013). *Study on performance evaluation of Sewage Treatment Plants (STPs) in Lucknow and Issue of wastewater sustainability*. <https://www.researchgate.net/publication/316515931>
- Dutta, V., Sharma, U., Iqbal, K., Adeeba, Kumar, R., & Pathak, A. K. (2018a). Impact of river channelization and riverfront development on fluvial habitat: evidence from Gomti River, a tributary of Ganges, India. *Environmental Sustainability*, 1(2), 167–184. <https://doi.org/10.1007/s42398-018-0016-0>
- Dutta, V., Sharma, U., Iqbal, K., Adeeba, Kumar, R., & Pathak, A. K. (2018b). Impact of river channelization and riverfront development on fluvial habitat: evidence from Gomti River, a tributary of Ganges, India. *Environmental Sustainability*, 1(2), 167–184. <https://doi.org/10.1007/s42398-018-0016-0>
- Ejiohuo, O., Onyeaka, H., Akinsemolu, A., Nwabor, O. F., Siyanbola, K. F., Tamasiga, P., & Al-Sharify, Z. T. (2025). Ensuring water purity: Mitigating environmental risks and safeguarding human health. *Water Biology and Security*, 4(2), 100341. <https://doi.org/10.1016/j.watbs.2024.100341>
- Goel, P., Saxena, A., Singh, D. Sen, & Verma, D. (2018). Impact of Rapid Urbanization on Water Quality Index in Groundwater Fed Gomati River, Lucknow, India. *Current Science*, 114(03), 650. <https://doi.org/10.18520/cs/v114/i03/650-654>
- Goumenou, M., & Tsatsakis, A. (2019). Proposing new approaches for the risk characterisation of single chemicals and chemical mixtures: The source related Hazard Quotient (HQS) and Hazard Index (HIS) and the adversity specific Hazard Index (HIA). *Toxicology Reports*, 6, 632–636. <https://doi.org/10.1016/j.toxrep.2019.06.010>
- Hughes, J., Cowper-Heays, K., Olesson, E., Bell, R., & Stroombergen, A. (2021). Impacts and implications of climate change on wastewater systems: A New Zealand perspective. *Climate Risk Management*, 31, 100262. <https://doi.org/10.1016/j.crm.2020.100262>
- Javan, K., Darestani, M., Ibrar, I., & Pignatta, G. (2025). Interrelated issues within the Water-Energy-Food nexus with a focus on environmental pollution for sustainable development: A review. *Environmental Pollution*, 368, 125706. <https://doi.org/10.1016/j.envpol.2025.125706>
- Kesari, K. K., Soni, R., Jamal, Q. M. S., Tripathi, P., Lal, J. A., Jha, N. K., Siddiqui, M. H., Kumar, P., Tripathi, V., & Ruokolainen, J. (2021). Wastewater Treatment and Reuse: a Review of its Applications and Health Implications. *Water, Air, & Soil Pollution*, 232(5), 208. <https://doi.org/10.1007/s11270-021-05154-8>
- Khan, R., Saxena, A., & Shukla, S. (2020). Evaluation of heavy metal pollution for River Gomti, in parts of Ganga Alluvial Plain, India. *SN Applied Sciences*, 2(8), 1451. <https://doi.org/10.1007/s42452-020-03233-9>
- Kumar, P. (2018a). Simulation of Gomti River (Lucknow City, India) future water quality under different mitigation strategies. *Heliyon*, 4(12), e01074. <https://doi.org/10.1016/j.heliyon.2018.e01074>
- Kumar, P. (2018b). Simulation of Gomti River (Lucknow City, India) future water quality under different mitigation strategies. *Heliyon*, 4(12), e01074. <https://doi.org/10.1016/j.heliyon.2018.e01074>
- Kumar, P., Srivastava, S., Banerjee, A., & Banerjee, S. (2022). Prevalence and predictors of water-borne diseases among elderly people in India: evidence from Longitudinal Ageing Study in India, 2017–18. *BMC Public Health*, 22(1), 993. <https://doi.org/10.1186/s12889-022-13376-6>
- Kumar Singh, P., & Verma, S. (2025). Water Pollution Analysis of Lucknow City: Sources, Impacts, and Mitigation Strategies. In *International Journal of Research Publication and Reviews Journal homepage: www.ijrpr.com* (Issue 6). www.ijrpr.com
- Landrigan, P. J., Stegeman, J. J., Fleming, L. E., Allemand, D., Anderson, D. M., Backer, L. C., Brucker-Davis, F., Chevalier, N., Corra, L., Czerucka, D., Bottein, M.-Y. D., Demeneix, B., Depledge, M., Deheyn, D. D., Dorman, C. J., Fénichel, P., Fisher, S., Gaill, F., Galgani, F., ... Rampal, P. (2020). Human Health and Ocean Pollution. *Annals of Global Health*, 86(1), 151. <https://doi.org/10.5334/aogh.2831>
- Lohani, M. B., Singh, A., Rupainwar, D. C., & Dhar, D. N. (2008). Seasonal variations of heavy metal contamination in river Gomti of Lucknow city region. *Environmental Monitoring and Assessment*, 147(1–3), 253–263. <https://doi.org/10.1007/s10661-007-0117-1>
- Mishra, A. K., Singh, J., & Mishra, P. P. (2024). *Microplastics in Freshwater Environment: A Case Study of River Gomti, Lucknow, India*. <https://doi.org/10.21203/rs.3.rs-5250209/v1>

- Mondal, S., & Patel, P. P. (2022). Incorporating Hydromorphological Assessments in the Fluvial Geomorphology Domain for Transitioning Towards Restorative River Science—Context, Concepts and Criteria. In *Fluvial Systems in the Anthropocene* (pp. 43–75). Springer International Publishing. https://doi.org/10.1007/978-3-031-11181-5_4
- Partani, S., Danandeh Mehr, A., Bostanmaneshrad, F., Arzhang, A., Pourrostami Niavol, K., & Nachtnebel, H.-P. (2024). Determining the main driver of hypoxia potential in freshwater inland lakes. *Journal of Cleaner Production*, 458, 142521. <https://doi.org/10.1016/j.jclepro.2024.142521>
- Parveen, N., & Singh, S. K. (2016a). Physico-Chemical Characteristics of Gomti River at Lucknow Stretch, India. *International Journal of Advanced Research*, 4(7), 749–759. <https://doi.org/10.21474/IJAR01/1163>
- Parveen, N., & Singh, S. K. (2016b). Physico-Chemical Characteristics of Gomti River at Lucknow Stretch, India. *International Journal of Advanced Research*, 4(7), 749–759. <https://doi.org/10.21474/IJAR01/1163>
- Paul, D. (2017). Research on heavy metal pollution of river Ganga: A review. *Annals of Agrarian Science*, 15(2), 278–286. <https://doi.org/10.1016/j.aasci.2017.04.001>
- Pericherla, S., & Vara, S. (2023). Effects of Diffused Pollution on Surface Waters Due to Agricultural Runoff. *Journal of Advanced Zoology*. <https://jzindia.com>
- Ravichandran, S., & Yadav, P. (2021). An overview of water pollution studies on Gomti river. *International Journal of Clinical Biochemistry and Research*, 8(2), 96–99. <https://doi.org/10.18231/j.ijcbr.2021.021>
- Ravindra Swaroop Sinha. (2021). *State of Groundwater in Uttar Pradesh A Situation Analysis with Critical Overview and Sustainable Solutions Ground Water Action Group* +.
- Rishita Jain and Khushi Dixit. (2023, July 1). *Namami Gange Programme: Projection vs Processing*. Global Association of Economics Education's GAEE.
- Siddig, M. M. S., Brevik, E. C., & Sauer, D. (2025). Human health risk assessment from potentially toxic elements in the soils of Sudan: A meta-analysis. *Science of The Total Environment*, 958, 178196. <https://doi.org/10.1016/j.scitotenv.2024.178196>
- Singh, B. J., Chakraborty, A., & Sehgal, R. (2023). A systematic review of industrial wastewater management: Evaluating challenges and enablers. *Journal of Environmental Management*, 348, 119230. <https://doi.org/10.1016/j.jenvman.2023.119230>
- Singh, H., Nielsen, M., & Greatrex, H. (2023). Causes, impacts, and mitigation strategies of urban pluvial floods in India: A systematic review. *International Journal of Disaster Risk Reduction*, 93, 103751. <https://doi.org/10.1016/j.ijdr.2023.103751>
- Singh, N., & Tung, S. (2025). Assessment of Water Quality of Gomti River at Lucknow. *Air, Soil and Water Research*, 18. <https://doi.org/10.1177/11786221251328589>
- Verma, R. (2025). *International Journal of Sciences and Innovation Engineering Microplastics: Environmental Ubiquity, Biological Fate, and Human Health Implications*. 2(8). <https://doi.org/10.70849/ijsci>
- Verma, R., & Choudhary, M. (2025). Utilizing Plastic Trash to Produce Char & Activated Carbon for Wastewater Treatment: A Review. *International Journal for Research in Applied Science and Engineering Technology*, 13(3), 1795–1798. <https://doi.org/10.22214/ijraset.2025.67664>
- Vetrimurugan, E., Brindha, K., Elango, L., & Ndwandwe, O. M. (2017). Human exposure risk to heavy metals through groundwater used for drinking in an intensively irrigated river delta. *Applied Water Science*, 7(6), 3267–3280. <https://doi.org/10.1007/s13201-016-0472-6>
- Wear, S. L., Acuña, V., McDonald, R., & Font, C. (2021). Sewage pollution, declining ecosystem health, and cross-sector collaboration. *Biological Conservation*, 255, 109010. <https://doi.org/10.1016/j.biocon.2021.109010>
- Weaver, D., Moyle, B. D., McLennan, C., & Casali, L. (2023). Taming the wicked problem of climate change with “virtuous challenges”: An integrated management heuristic. *Journal of Environmental Management*, 347, 119136. <https://doi.org/10.1016/j.jenvman.2023.119136>
- Yaduvanshi, N., Ghildiyal, D., Pandey, R., & . Y. (2024). Water Quality Parameters of Major Rivers of Uttar Pradesh in Winter Season. *International Journal of Lakes and Rivers*, 17(1), 29–46. <https://doi.org/10.37622/ijlr/17.1.2024.29-46>